

LA-UR-26-27391

Approved for public release; distribution is unlimited.

Title: A Compact ENDF (ACE) Format Specification

Author(s): Conlin, Jeremy Lloyd
Haeck, Wim
Romano, Paul

Intended for: Report

Issued: 2026-08-14



Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

A Compact ENDF (ACE) Format Specification

Jeremy Lloyd Conlin (editor)

Los Alamos National Laboratory

Contributors:

Jeremy Lloyd Conlin (*Los Alamos National Laboratory*)

Wim Haeck (*Los Alamos National Laboratory*)

Paul Romano (*Argonne National Laboratory*)

Contents

1	Introduction	1
1.1	Types of ACE-Formatted Data	1
1.2	Classes of ACE-Formatted Data	1
1.3	ACE Libraries	1
2	ACE Tables	2
2.1	ACE Header	2
2.1.1	IZAW Array	4
2.1.2	NXS Array	4
2.1.3	JXS Array	4
2.2	The XSS Array	4
3	Unique ACE Table Identifier	6
3.1	ZAID	6
3.2	SZAID	6
4	Continuous-Energy and Discrete Neutron Transport Tables	7
4.1	NXS Array	10
4.2	JXS Array	11
4.3	Format of Individual Data Blocks	12
4.3.1	ESZ Block	12
4.3.2	NU and DNU Blocks	12
4.3.3	BDD Block	13
4.3.4	MTR, MTRP and MTRH Blocks	14
4.3.5	LQR Block	15
4.3.6	TYR and TYRH Blocks	16
4.3.7	LSIG, LSIGP and LSIGH Blocks	17
4.3.8	SIG Block	17
4.3.9	LAND, LANDP and LANDH Blocks	18
4.3.10	AND, ANDP and ANDH Blocks	19
4.3.11	LDLW, LDLWP, DNEDL and LDLWH Blocks	21
4.3.12	DLW, DLWP, DLWH and DNED Blocks	22
4.3.12.1	LAW=1—Tabular Equiprobable Energy Bins	24
4.3.12.2	LAW=2—Discrete Photon Energy	25
4.3.12.3	LAW=3—Level Scattering	25
4.3.12.4	LAW=4—Continuous Tabular Distribution	26
4.3.12.5	LAW=5—General Evaporation Spectrum	27

4.3.12.6	LAW=7—Simple Maxwellian Fission Spectrum	27
4.3.12.7	LAW=9—Evaporation Spectrum	28
4.3.12.8	LAW=11—Energy Dependent Watt Spectrum	29
4.3.12.9	LAW=22—Tabular Linear Functions of Incident Energy Out	30
4.3.12.10	LAW=24—Tabular Energy Multipliers	31
4.3.12.11	LAW=44—Kalbach-87 Formalism	32
4.3.12.12	LAW=61—Like LAW=44, but tabular angular distribution instead of Kalbach-87	33
4.3.12.13	LAW=66— N -body phase space distribution	34
4.3.12.14	LAW=67—Laboratory Angle-Energy Law	35
4.3.12.15	Energy-Dependent Neutron Yields	36
4.3.13	GPD Block	37
4.3.14	SIGP and SIGH Blocks	38
4.3.15	YP and YH Blocks	40
4.3.16	FIS Block	41
4.3.17	UNR Block	41
4.3.18	PTYPE Block	43
4.3.19	NTRO Block	44
4.3.20	IXS Block	44
4.3.21	HPD Block	45
5	Neutron Dosimetry	46
5.1	NXS Array	46
5.2	JXS Array	46
6	Thermal Scattering	47
6.1	NXS Array	47
6.2	JXS Array	47
6.3	Format of Individual Data Blocks	48
6.3.1	ITIE Block	48
6.3.2	ITXE Block	49
6.3.3	ITCE Block	51
6.3.4	ITCA Block	52

7	Continuous-Energy Photon	53
7.1	NXS Array	53
7.2	JXS Array	53

Todo list

This needs to be done. 6

1 Introduction

The ACE format consists of two *types* and many *classes* of data. The data are kept in an ACE Table. The term ACE Table and ACE file are often used interchangeably.

1.1 Types of ACE-Formatted Data

There are two types of ACE-formatted data; simply called Type 1 and Type 2.

Type 1 Standard formatted tables. These tables contain ASCII text and are machine independent; they are readable on every machine.

Type 2 Standard unformatted tables. These tables are binary and can be generated from the Type 1 files. They are more compact and faster to read than the Type 1 ACE Tables but are machine/platform dependent; they are not readable on every machine.

Traditionally Type 2 ACE files were more commonly used because they were smaller in size and faster to read. However due to the fact that they are not portable across machines and platforms they have fallen out of fashion.

1.2 Classes of ACE-Formatted Data

There are many classes of ACE-formatted data:

1. continuous-energy neutron (see Section 4),
2. discrete-reaction neutron,
3. neutron dosimetry (see Section 5),
4. $S(\alpha, \beta)$ thermal scattering (see Section 6),
5. continuous-energy photoatomic (see Section 7),
6. continuous-energy electron interaction,
7. continuous-energy photonuclear interaction,
8. multigroup-energy neutron, and
9. multigroup-energy photoatomic.

Each of these classes of data are described later in this document.

An ACE Table is an entity that contains evaluation-dependent data about one of the many classes of data for a specific material—an target isotope, isomer, or element. For a given ZAID, the data contained on a Type 1 and Type 2 tables are identical. Simulations run with one type of data should produce identical results as those run with the other type of data.

1.3 ACE Libraries

A collection of ACE data tables that derive from a single set of evaluation files are typically grouped together in a “library”—not to be confused from the evaluation library from which they derive. Multiple ACE data tables can concatenated into the same logical file on the computer, although this has fallen somewhat out of fashion due to the large amount of data on each ACE table derived from modern evaluation files. Applications that use ACE-formatted data should produce the same results regardless of whether the tables are contained in one logical file on the computer or spread across many.

2 ACE Tables

An ACE Table consists of a Header followed by an array (XSS) containing the actual data. The Header and XSS array are the same regardless of whether the ACE Table is Type 1 or Type 2. Each line in a Type 1 ACE Table is 80 characters or less.

2.1 ACE Header

The first section of an ACE Table is the Header. The ACE Header contains metadata¹ about the ACE Table. The Header consists of four parts:

1. Opening,
2. IZAW array,
3. NXS array, and
4. JXS array.

An example of an ACE Table Header (from ¹H in the ENDF71x library) is given in Figure 1 with each part highlighted a different color.

1	1001.80c	0.999167	2.5301E-08	12/17/12				
2	H1 ENDF71x (jlconlin)	Ref. see jlconlin	(ref 09/10/2012	10:00:53)			mat 125	
3	0	0.	0	0.	0	0.	0	0.
4	0	0.	0	0.	0	0.	0	0.
5	0	0.	0	0.	0	0.	0	0.
6	0	0.	0	0.	0	0.	0	0.
7	17969	1001	590	3	0	1	1	0
8	0	1	1	0	0	0	0	0
9	1	0	2951	2954	2957	2960	2963	4352
10	4353	5644	5644	5644	6234	6235	6236	6244
11	6245	6245	6246	16721	0	16722	0	0
12	0	0	0	0	0	16723	16724	16725

Figure 1: Header example. The (Legacy) Opening (lines 1–2) is in red, the IZAW array (lines 3–6) is in blue, the NXS array (lines 7–8) is in teal, and the JXS array (lines 9–12) is in violet.

Legacy Header Opening There are two slightly different formats for the Header Opening. The most common one found is called here the Legacy Opening and is the one demonstrated in the Header example in Figure 1.

The Legacy Opening consists of several variables given over two 80-character lines. The variables and the Fortran format for reading the variable is given in Table 1

2.0.1 Header Opening There is a limitation to the number of unique ZA IDs for a given ZA; 100 different IDs, in fact, for each class of ACE Table. To overcome this limitation, a new Header

¹data about the data

Line	Variable	Format	Description
1	HZ	A10	ZAID (see Section 3.1)
1	AW	E12.0	Atomic weight ratio
1	TZ	E12.0	Temperature
1	—	1X	(blank space)
1	HD	A10	Processing date
2	HK	A70	Descriptive string
2	HM	A10	10-character material identifier

Table 1: Variables in the Legacy Opening part of the ACE Header.

Line	Variable	Format	Description
1	VERS	A10	Version format string
1	—	1X	(blank space)
1	HZ	A24	SZAID (see Section 3.2)
1	—	1X	(blank space)
1	SRC	A24	Evaluation source
2	AW	E12.0	Atomic weight ratio
2	—	1X	(blank space)
2	TZ	E12.0	Temperature
2	—	1X	(blank space)
2	HD	A10	Processing date
2	—	1X	(blank space)
2	N	I4	Number of comment lines to follow
3-(N+2)	—	A70	N comment lines

Table 2: Variables in the 2.0.1 Opening part of the ACE Header.

Opening[Conlin:2012Updat-0] was developed in 2012 and updated a few years later to correct some errors.

2.0.1	1001.800nc	ENDF/B-VIII.0-B1
0.999167	2.5301e-08	2018-05-02 2
1001.00c	0.999167	2.5301E-08 05/02/18
H1 Lib80x (jlconlin)	Ref. see jlconlin (ref 01/29/2018 07:54)	mat 125

Figure 2: Header Opening example. The Legacy Opening is shown in blue while the 2.0.1 Opening consists of the red and the blue portions.

Note that a Legacy Header Opening can be contained in the comment section of the 2.0.1 Header

Opening. This was designed explicitly to allow backwards compatibility while application codes were modified to be able to handle. An example of this is shown in Figure 2. Codes that cannot read the 2.0.1 Header can be told (typically via an `xmdir[Conlin:2012Updat-0]` entry) to start reading the ACE Table several lines after the beginning of the 2.0.1 Header.

Following the Opening of the Header are three arrays, IZAW, NXS, and JXS respectively. They are each described below. Immediately following the JXS array is the XSSarray.

2.1.1 IZAW Array

The IZAW array follows on the lines immediately following the Header. It consists of 16 pairs of ZA's (IZ) and atomic weight ratios (AW). The IZ entries are still needed for $S(\alpha, \beta)$ Tables to indicate for which isotope(s) the scattering data are appropriate.

The 16 pairs of numbers are spread over 4 lines. The Fortran format for reading/writing the numbers on one line is: `4(I7,F11.0)`.

2.1.2 NXS Array

The NXS array comes on the 2 lines after the IZAW array. The NXS array has 16 integer elements; 8 on each line. The Fortran format for reading/writing the numbers on each line is: `8I9`. The first element of the NXS array indicates how many numbers are in the XSS array. The remainder of the NXS array elements (usually) indicate how many of different pieces of data there is.

2.1.3 JXS Array

The JXS array comes on the 4 lines after the NXS array. The JXS array has 32 integer elements; 8 on each line. The Fortran format for reading/writing the numbers on each line is: `8I9`. The JXS array contains indices to the XSS array where different pieces of data begins.

The specific definition of the elements of the NXS and JXS arrays are dependent on the class of data in the Table and are defined in the section of this document that describes each class of data.² Note that not all elements of the arrays are (currently) being used, allowing for future expansion.

2.2 The XSS Array

After the ACE Header comes the XSS array. It is typically *very* large with hundreds of thousands of elements. It is broken up into blocks with the blocks being dependent on the class of data that is contained in the table. The description and definition of each of these blocks can be found in the descriptions later in this document.

The data is written with 4 floating-point numbers on each 80-character line. All data in the XSS array can be read using the Fortran format: `4E20.0` for each line.

²See, for example, Table 3 and Table 4.

```

2.0.1                1001.710nc                ENDFB-VII.1
  0.999167    2.5301E-08    12/17/12    3
The next two lines are the first two lines of 'old-style' ACE.
1001.80c    0.999167    2.5301E-08    12/17/12
H1 ENDF71x (jlconlin) Ref. see jlconlin (ref 09/10/2012 10:00:53) mat 125
  0          0.          0          0.          0          0.          0          0.
  0          0.          0          0.          0          0.          0          0.
  0          0.          0          0.          0          0.          0          0.
  0          0.          0          0.          0          0.          0          0.
17969    1001    590          3          0          1          1          0
  0          1          1          0          0          0          0          0
  1          0    2951    2954    2957    2960    2963    4352
4353    5644    5644    5644    6234    6235    6236    6244
6245    6245    6246    16721          0    16722          0          0
  0          0          0          0          0    16723    16724    16725
1.00000000000E-11  1.03125000000E-11  1.06250000000E-11  1.09375000000E-11
1.12500000000E-11  1.15625000000E-11  1.18750000000E-11  1.21875000000E-11
1.25000000000E-11  1.28125000000E-11  1.31250000000E-11  1.34375000000E-11
1.37500000000E-11  1.43750000000E-11  1.50000000000E-11  1.56250000000E-11
1.62500000000E-11  1.68750000000E-11  1.75000000000E-11  1.81250000000E-11
1.87500000000E-11  1.93750000000E-11  2.00000000000E-11  2.09375000000E-11
2.18750000000E-11  2.28125000000E-11  2.37500000000E-11  2.46875000000E-11
2.56250000000E-11  2.65625000000E-11  2.75000000000E-11  2.84375000000E-11
2.93750000000E-11  3.03125000000E-11  3.12500000000E-11  3.21875000000E-11
3.31250000000E-11  3.40625000000E-11  3.50000000000E-11  3.59375000000E-11

```

Figure 3: ACE Header with beginning of XSS array for ^1H . Note this uses the 2.0.1 Header with backwards compatibility with the Legacy Header.

3 Unique ACE Table Identifier

This needs to be done.

Each ACE Table needs to have an identifier to uniquely distinguish the data that is contained in the Table.

3.1 ZAID

3.2 SZAID

With the introduction of the 2.0.1 ACE Header, the identifier was modified to better specify the metastable state of the material as well as expand the available space for identifiers.

The new identifier is referred to as a SZAID³.

³pronounced “ess-ZAID”

4 Continuous-Energy and Discrete Neutron Transport Tables

The format of individual blocks found on neutron transport tables is identical for continuous-energy and discrete-reaction ACE Tables; the format for both are described in this section. The blocks of data are:

1. **ESZ Block**—contains the main energy grid for the Table and the total, absorption, and elastic cross sections as well as the average heating numbers. The **ESZ Block** always exists. See Section 4.3.1.
2. **NU Block**—contains prompt, delayed and/or total $\bar{\nu}$ as a function of incident neutron energy. The **NU Block** exists only for fissionable isotopes; that is, if $JXS(2) \neq 0$. See Section 4.3.2.
3. **MTR Block**—contains a list of ENDF MT numbers for all neutron reactions other than elastic scattering. The **MTR Block** exists for all isotopes that have reactions other than elastic scattering; that is, all isotopes with $NXS(4) \neq 0$. See Section 4.3.4.
4. **LQR Block**—contains a list of kinematic Q -values for all neutron reactions other than elastic scattering. The **LQR Block** exists if $NXS(4) \neq 0$. See Section 4.3.5.
5. **TYR Block**—contains information about the type of reaction for all neutron reactions other than elastic scattering. Information for each reaction includes the number of secondary neutrons and whether secondary neutron angular distributions are in the laboratory or center-of-mass system. The **TYR Block** exists if $NXS(4) \neq 0$. See Section 4.3.6.
6. **LSIG Block**—contains a list of cross section locators for all neutron reactions other than elastic scattering. The **LSIG Block** exists if $NXS(4) \neq 0$. See Section 4.3.7.
7. **SIG Block**—contains cross sections for all reactions other than elastic scattering. The **SIG Block** exists if $NXS(4) \neq 0$. See Section 4.3.8.
8. **LAND Block**—contains a list of angular-distribution locators for all reactions producing secondary neutrons. The **LAND Block** always exists. See Section 4.3.9.
9. **AND Block**—contains list angular distributions for all reactions producing secondary neutrons. The **AND Block** always exists. See Section 4.3.10.
10. **LDLW Block**—contains a list of energy distributions for all reactions producing secondary neutrons except for elastic scattering. The **LDLW Block** exists if $NXS(5) \neq 0$. See Section 4.3.11.
11. **DLW Block**—contains energy distributions for all reactions producing secondary neutrons except for elastic scattering. The **DLW Block** exists if $NXS(5) \neq 0$. See Section 4.3.12.
12. **GPB Block**—contains the total photon production cross section tabulated on the ESZ energy grid and a $30 \times$ matrix of secondary photon energies. The **GPB Block** exists only for those older evaluations that provide coupled neutron/photon information; that is, if $JXS(12) \neq 0$. See Section 4.3.13.
13. **MTRP Block**—contains a list of MT numbers for all photon production reactions. The term “photon production reaction” is used for any information describing a specific neutron-in, photon-out reaction. The **MTRP Block** exists if $NXS(6) \neq 0$. See Section 4.3.4.
14. **LSIGP Block**—contains a list of cross section locators for all photon production reactions. The **LSIGP Block** exists if $NXS(6) \neq 0$. See Section 4.3.7.
15. **SIGP Block**—contains cross sections for all photon production reactions. The **SIGP Block**

- exists if $\text{NXS}(6) \neq 0$. See Section 4.3.14.
16. **LANDP Block**—contains a list of angular-distribution locators for all photon production reactions. The **LANDP Block** exist if $\text{NXS}(6) \neq 0$. See Section 4.3.9
 17. **ANDP Block**—contains photon angular distributions for all photon production reactions. The **ANDP Block** exists if $\text{NXS}(6) \neq 0$. See Section 4.3.10.
 18. **LDLWP Block**—contains a list of energy-distribution locators for all photon production reactions. The **LDLWP Block** exists if $\text{NXS}(6) \neq 0$. See Section 4.3.11.
 19. **DLWP Block**—contains photon energy distributions for all photon production reactions. The **DLWP Block** exists if $\text{NXS}(6) \neq 0$. See Section 4.3.12.
 20. **YP Block**—contains a list of MT identifiers of neutron reaction cross sections required as photon production yield multipliers. The **YP Block** exists if $\text{NXS}(6) \neq 0$. See Section 4.3.15.
 21. **FIS Block**—contains the total fission cross section tabulated on the ESZ energy grid. The **FIS Block** exists if $\text{JXS}(21) \neq 0$. See Section 4.3.16.
 22. **UNR Block**—contains the unresolved resonance range probability tables. The **UNR Block** exists if $\text{JXS}(23) \neq 0$. See Section 4.3.17.
 23. **PTYPE Block**—contains a list of particle types for which production data will be given. The **PTYPE Block** exists if $\text{JXS}(30) \neq 0$. See Section 4.3.18.
 24. **NTRO Block**—contains the number of reactions that produce the corresponding particle type given in the **PTYPE Block**. The **NTRO Block** exists if $\text{JXS}(31) \neq 0$. See Section 4.3.19.
 25. **IXS Block**—particle production data locators for each particle type given in the **PTYPE Block**. The **IXS Block** exists if $\text{JXS}(32) \neq 0$. See Section 4.3.20.
 26. **HPD Block**—total particle production cross section and average heating numbers for the current particle type. The **HPD Block** for a given particle type i exists if $\text{JXS}(32) + 10 * (i - 1) \neq 0$. See Section 4.3.21.
 27. **MTRH Block**—contains a list of ENDF MT numbers for all reactions that produce the current particle. The **MTR Block** for a given particle type i exists if $\text{JXS}(32) + 10 * (i - 1) + 1 \neq 0$. See Section 4.3.4.
 28. **TYRH Block**—contains the reaction types for all reactions that produce the current particle. The **TYRH Block** for a given particle type i exists if $\text{JXS}(32) + 10 * (i - 1) + 2 \neq 0$. See Section 4.3.6.
 29. **LSIGH Block**—contains the cross section locators for all reactions that produce the current particle. The **LSIGH Block** for a given particle type i exists if $\text{JXS}(32) + 10 * (i - 1) + 3 \neq 0$. See Section 4.3.7.
 30. **SIGH Block**—contains the cross section data for all reactions that produce the current particle. The **SIGH Block** for a given particle type i exists if $\text{JXS}(32) + 10 * (i - 1) + 4 \neq 0$. See Section 4.3.14.
 31. **LANDH Block**—contains the angular distribution locators for all reactions that produce the current particle. The **LANDH Block** for a given particle type i exists if $\text{JXS}(32) + 10 * (i - 1) + 5 \neq 0$. See Section 4.3.9.
 32. **ANDH Block**—contains the angular distribution data for all reactions that produce the current particle. The **ANDH Block** for a given particle type i exists if $\text{JXS}(32) + 10 * (i - 1) + 6 \neq 0$. See Section 4.3.10.

33. **LDLWH Block**—contains the energy distribution locators for all reactions that produce the current particle. The **LDLWH Block** for a given particle type i exists if $JXS(32) + 10 * (i - 1) + 7 \neq 0$. See Section 4.3.11.
34. **DLWH Block**—contains the energy distribution data for all reactions that produce the current particle. The **DLWH Block** for a given particle type i exists if $JXS(32) + 10 * (i - 1) + 8 \neq 0$. See Section 4.3.12.
35. **YH Block**—contains the particle production yield multiplier for all reactions that produce the current particle. The **YH Block** for a given particle type i exists if $JXS(32) + 10 * (i - 1) + 9 \neq 0$. See Section 4.3.12.

4.1 NXS Array

Table 3: NXS array element definitions for NXS ACE Table.

Element	Name	Description
1	—	Length of second block of data (XSS array)
2	ZA	$1000 * Z + A$
3	NES	Number of energies
4	NTR	Number of reactions excluding elastic scattering
5	NR	Number of reactions having secondary neutrons excluding elastic scattering
6	NTRP	Number of photon production reactions
7	NTYPE	Number of particle types for which production data is given
8	NPCR	Number of delayed neutron precursor families
9	S	Excited state [†]
10	Z	Atomic number [†]
11	A	Atomic mass number [†]
	...	
14		Reserved [‡]
15		Reserved [‡]
16		Reserved [‡]

[†] These values were introduced with the new 2.0.0 Header[Conlin:2012Updat-0].

[‡] These entries are reserved for the use of transport codes (i.e., MCNP).

4.2 JXS Array

Table 4: JXS array element definitions for JXS ACE Table.

Element	Name	Location Description
1	ESZ	Energy table
2	NU	Fission ν data
3	MTR	MT array
4	LQR	Q -value array
5	TYR	Reaction type array
6	LSIG	Table of cross section locators
7	SIG	Cross sections
8	LAND	Table of angular distribution locators
9	AND	Angular distributions
10	LDLW	Table of energy distribution locators
11	DLW	Energy distributions
12	GPD	Photon production data
13	MTRP	Photon production MT array
14	LSIGP	Table of photon production cross section locators
15	SIGP	Photon production cross sections
16	LANDP	Table of photon production angular distribution locators
17	ANDP	Photon production angular distributions
18	LDLWP	Table of photon production energy distribution locators
19	DLWP	Photon production energy distributions
20	YP	Table of yield multipliers
21	FIS	Total fission cross section
22	END	Last word of the conventional table (last word of photon production data)
23	LUNR	Probability tables
24	DNU	Delayed $\bar{\nu}$ data
25	BDD	Basic delayed neutron precursor data (λ 's, probabilities)
26	DNEDL	Table of delayed neutron energy distribution locators
27	DNED	Delayed neutron energy distributions
	...	
30	PTYPE	Particle type array
31	NTRO	Array containing the number of particle production reactions
32	NEXT	Table of particle production locators (IXS array)

4.3 Format of Individual Data Blocks

4.3.1 ESZ Block

The **ESZ Block** provides the common incident energy table for all reactions defined in the ACE Table, cross section tables for fundamental cross sections (total, absorption, and elastic scattering) and average heating numbers. The format of the **ESZ Block** is given in Table 5. The starting index ESZ for this block is given by JXS(1).

Table 5: ESZ Block.

Location in XSS	Parameter	Description
S_{ESZ}	$E(l), l = 1, \dots, N_E$	Energies
$S_{\text{ESZ}} + N_E$	$\sigma_t(l), l = 1, \dots, N_E$	Total cross section
$S_{\text{ESZ}} + 2N_E$	$\sigma_a(l), l = 1, \dots, N_E$	Total neutron disappearance cross section [†]
$S_{\text{ESZ}} + 3N_E$	$\sigma_{el}(l), l = 1, \dots, N_E$	Elastic cross section
$S_{\text{ESZ}} + 4N_E$	$H_{ave}(l), l = 1, \dots, N_E$	Average Heating numbers

[†] The disappearance cross section is defined in [Trkov:2011ENDF-0] as MT101

Note: S_{ESZ} is index of the XSS array where the **ESZ Block** starts, JXS(1), and N_E is the number of energy energy points, NXS(3).

4.3.2 NU and DNU Blocks

The **NU Block** is used to specify prompt and/or total $\bar{\nu}$ and is present only if JXS(2) > 0. Delayed $\bar{\nu}$ data is specified in the **DNU Block** (which is only present if JXS(24) > 0) but it shares some of the tables defined in this section.

When it is present, there are two possibilities for the **NU Block**:

1. **Either prompt or total $\bar{\nu}$ is given (but not both).** ($\text{XSS}(\text{JXS}(2)) > 0$)
A single $\bar{\nu}$ array is given and it begins at location XSS(KNU) where $\text{KNU} = \text{JXS}(2)$.
2. **Both prompt and total $\bar{\nu}$ are given.** ($\text{XSS}(\text{JXS}(2)) < 0$). Two $\bar{\nu}$ arrays are given, one for prompt $\bar{\nu}$ and another for total $\bar{\nu}$. The absolute value of XSS(JXS(2)) is the location of the total $\bar{\nu}$ array so that the locations for the two $\bar{\nu}$ arrays are as follows:
 - The prompt $\bar{\nu}$ array begins at XSS(KNU) where $\text{KNU} = \text{JXS}(2) + 1$.
 - The total $\bar{\nu}$ array begins at XSS(KNU) where $\text{KNU} = \text{JXS}(2) + \text{ABS}(\text{XSS}(\text{JXS}(2))) + 1$.

There are two possible forms for these $\bar{\nu}$ arrays; either polynomial (see Table 6) or tabulated (see Table 7). The format is specified by the LNU flag located in the XSS array at index KNU where KNU is defined above.

Table 6: NU Block—Polynomial function form.

Location in XSS	Parameter	Description
KNU	LNU=1	Polynomial function flag
KNU+1	N_C	Number of coefficients
KNU+2	$C(l), l = 1, \dots, N_C$	Coefficients

When using the polynomial function form of the $\bar{\nu}$ array, $\bar{\nu}$ is reconstructed as

$$\bar{\nu}(E) = \sum_{l=1}^{N_C} C(l) E^{l-1}, \quad (1)$$

where the energy, E , is given in MeV.

Table 7: NU Block—Tabulated form.

Location in XSS	Parameter	Description
KNU	LNU=2	Tabulated data flag
KNU+1	N_R	Number of interpolation regions
KNU+2	$\text{NBT}(l), l = 1, \dots, N_R$	ENDF interpolation parameters
KNU+2+ N_R	$\text{INT}(l), l = 1, \dots, N_R$	ENDF interpolation scheme [†]
KNU+2+2 N_R	N_E	Number of energies
KNU+3+2 N_R	$E(l), l = 1, \dots, N_E$	Tabulated energy points
KNU+3+2 $N_R + N_E$	$\bar{\nu}(l), l = 1, \dots, N_E$	Tabulated $\bar{\nu}$ values

[†] If $N_R = 0$, NBT and INT are omitted and linear-linear interpolation is assumed.

For the [DNU Block](#), the delayed $\bar{\nu}$ array begins at `XSS(KNU)` where `KNU = JXS(24)`. Delayed $\bar{\nu}$ must be given in the tabulated form as described in Table 7. The polynomial form is not allowed in the [DNU Block](#).

4.3.3 BDD Block

The [BDD Block](#) is used to specify basic delayed neutron precursor data and is present only if `JXS(25) > 0`. For every precursor group (the total number of precursor groups is given in `NXS(8)`), a decay constant is given along with the partial probability that a delayed fission neutron is born from the current group. This data is given in the format given in table Table 8. The starting index BDD for this block is given by `JXS(25)`.

Table 8: Delayed $\bar{\nu}$ precursor distribution..

Location in XSS	Parameter	Description
Data for precursor group 1		
BDD	DEC_1	Decay constant for the group 1
BDD + 1	N_R	Number of interpolation regions
BDD + 2	$NBT(l), l = 1, \dots, N_R$	ENDF interpolation parameters [†]
BDD + 2 + N_R	$INT(l), l = 1, \dots, N_R$	ENDF interpolation scheme
BDD + 2 + $2N_R$	N_E	Number of energies
BDD + 3 + $2N_R$	$E(l), l = 1, \dots, N_E$	Tabulated energy points
BDD + 3 + $2N_R + N_E$	$P(l), l = 1, \dots, N_E$	Corresponding probabilities
Data for precursor group 2—same format as for group 1		
...		
Data for precursor group NPCR = NXS(8)—same format as for group 1		

[†] If $N_R = 0$, NBT and INT are omitted and linear-linear interpolation is assumed.

4.3.4 MTR, MTRP and MTRH Blocks

The format of the [MTR Block](#) (for incident neutron reactions), [MTRP Block](#) (for photon production reactions) and [MTRH Block](#) (for particle production reactions), is given in Table 10 and provides a list of MT numbers for which data is available in other blocks of the ACE Table. The starting index depends on whether it is the [MTR Block](#), [MTRP Block](#) or [MTRH Block](#) and are given in Table 9. For the particle production [MTRH Block](#), i refers to the index of the corresponding particle type defined on the [PTYPE Block](#) and is between 1 and NTYPE.

Block	LMT	NMT
MTR	JXS(3)	NXS(4)
MTRP	JXS(13)	NXS(6)
MTRH	XSS(JXS(32)+10*(i-1)+1)	XSS(JXS(31)+i-1)

 Table 9: LMT and NMT values for the [MTR Block](#) and [MTRH Block](#).

Table 10: MTR & MTRP Block.

Location in XSS	Parameter	Description
LMT	MT_1	First ENDF Reaction available
LMT+1	MT_2	Second ENDF Reaction available
...		

Continued on next page

Table 10: MTR & MTRP Block (continued)

Location in XSS	Parameter	Description
LMT+NMT-1	MT _{NMT}	Last ENDF reaction available

For the [MTR Block](#) and [MTRH Block](#), MT₁, ..., MT_{NMT} are standard ENDF MT numbers; that is, MT=16=(*n*, 2*n*); MT=17=(*n*, 3*n*); etc. For a complete listing of MT numbers, see [Trkov:2011ENDF-0]. It is important to note here that the order in which these MT numbers are given is not arbitrary. The first NXS(5) values will be the MT numbers of reactions that produce secondary particles of the same type as the incident particle (i.e. there is secondary particle distribution data for the incident particle type for these reactions). The next NXS(4) – NXS(5) values will then be the MT numbers for reactions that do not produce a secondary particle of the same type as the incident particle.

For the [MTR Block](#), every MT number may appear only once. In the [MTRH Block](#), it is possible for a given MT number to appear twice if the same particle is also produced as the residual after the reaction (e.g. d + t -> d + d).

For the [MTRP Block](#), the MT numbers are somewhat arbitrary. To understand the scheme used for numbering the photon production MTs, it is necessary to realize that in the ENDF format, more than one photon can be produced by a particular neutron reaction that is itself specified by a single MT. Each of these photons is produced with an individual energy-dependent cross section. For example, MT102 (radiative capture) might be responsible for 40 photons, each with its own cross section, angular distribution, and energy distribution. We need 40 photon MTs to represent the data; the MTs are numbered 102001, 102002, ..., 102040. Therefore, if ENDF MT *N* is responsible for *M* photons, we shall number the photon MTs 1000**N*+1, 1000**N*+2, ..., 1000**N*+*M*.

4.3.5 LQR Block

The format of the [LQR Block](#), containing the reaction-specific *Q*-values, is given in Table 11. The index at the start of the [LQR Block](#), *S*_{LQR}=JXS(4). The number of reactions, NMT, is the same through the ACE Table, NMT=NXS(4).

Table 11: LQR Block.

Location in XSS	Parameter	Description
<i>S</i> _{LQR}	<i>Q</i> ₁	<i>Q</i> -value for reaction MT ₁
<i>S</i> _{LQR} +1	<i>Q</i> ₂	<i>Q</i> -value for reaction MT ₂
...		
<i>S</i> _{LQR} +NMT-1	<i>Q</i> _{NMT}	<i>Q</i> -value for reaction MT _{NMT}

4.3.6 TYR and TYRH Blocks

The format of the [TYR Block](#) (for incident neutron reactions) and [TYRH Block](#) (for particle production reactions) is given in Table 13. The starting index LTYR depends on whether it is the [TYR Block](#) or [TYRH Block](#) and is given in Table 12. For the particle production [TYRH Block](#), i refers to the index of the corresponding particle type defined on the [PTYPE Block](#) and is between 1 and NTYPE.

Block	LTYR	NMT
TYR	JXS(5)	NXS(4)
TYRH	XSS(JXS(32)+10*(i-1)+2)	XSS(JXS(31)+i-1)

Table 12: LTYR and NMT values for the [TYR Block](#) and [TYRH Block](#).

Table 13: TYR Block.

Location in XSS	Parameter	Description
S_{LTYR}	TY_1	Particle release for reaction MT_1
$S_{\text{LTYR}+1}$	TY_2	Particle release for reaction MT_2
...		
$S_{\text{LTYR}+\text{NMT}-1}$	TY_{NMT}	Particle release for reaction MT_{NMT}

The possible values of TY are $\pm 1, \pm 2, \pm 3, \pm 4, \pm 19, 0$, and integers greater than 100 in absolute value; the sign indicates the system for scattering:

negative center-of-mass,

positive Lab.

Thus if $\text{TY}_i = +3$, three particles are released for reaction MT_i and the data on the cross section tables used to determine the exiting neutrons' angles are given in the Lab frame of reference. $\text{TY} = 19$ indicates fission (only used in the [TYR Block](#)). The number of secondary neutrons released is determined from the fission $\bar{\nu}$ data found in the [NU Block](#). $\text{TY}_i = 0$ indicates absorption; no particles are released. $|\text{TY}_i| > 100$ signifies reactions other than fission that have energy-dependent multiplicities (currently only used in the [TYR Block](#)). The number of secondary particles released is determined from the yield data found in the [DLW Block](#) or [DLWH Block](#). The MT_i s are given in the [MTR Block](#) or [MTRH Block](#).

As the elastic scattering reaction is not included in the [TYR Block](#), the reference frame used for this reaction is not given in the [TYR Block](#) nor anywhere else in the ACE file. The reference frame for elastic scattering is always assumed to be the center-of-mass system, since this is the way the data has to be given in the ENDF evaluation [Trkov:2011ENDF-0].

4.3.7 LSIG, LSIGP and LSIGH Blocks

The **LSIG Block** (for incident neutron cross sections), **LSIGP Block** (for photon production cross sections) and **LSIGH Block** (for particle production cross sections), give the locators for the cross section array for each reaction MT. A locator is a *relative* index in the XSS array where some piece of data can be found. In this case, the data are the cross section values. The format of the **LSIG Block**, **LSIGP Block** and **LSIGH Block** is given in Table 15. The format for the incident neutron cross section arrays is given in Section 4.3.8, the format for the photon production cross sections is given in Section 4.3.14 and the format for the particle production cross sections is given in Section 4.3.14.

The starting index LXS depends on whether it is the **LSIG Block**, **LSIGP Block** or **LSIGH Block** and are given in Table 14. For the particle production **LSIGH Block**, i refers to the index of the corresponding particle type defined on the **PTYPE Block** and is between 1 and NTYPE.

Block	LXS	NMT
LSIG	JXS(6)	NXS(4)
LSIGP	JXS(14)	NXS(6)
LSIGH	XSS(JXS(32)+10*(i-1)+3)	XSS(JXS(31)+i-1)

Table 14: TYR and NMT values for the **TYR Block** and **TYRH Block**.

The MTs are given in the **MTR Block**, the **MTRP Block** and **MTRH Block** respectively for the **LSIG Block**, the **LSIGP Block** and **LSIGH Block** respectively. LOCA_i must be monotonically increasing. All locators (LOCA) are *relative* to $\text{SIG}=\text{JXS}(7)$ for the **LSIG Block**, *relative* to $\text{SIGP}=\text{JXS}(14)$ for the **LSIGP Block** and *relative* to $\text{ANDH}=\text{XSS}(\text{JXS}(32)+10*(i-1)+4)$ for the **LSIGH Block** for particle index i .

Table 15: LSIG & LSIGP Block.

Location in XSS	Parameter	Description
LXS	LOCA_1	Location of cross sections for reaction MT_1
LXS+1	LOCA_2	Location of cross sections for reaction MT_2
...		
LXS+NMT-1	LOCA_{NMT}	Location of cross sections for reaction MT_{NMT}

4.3.8 SIG Block

The **SIG Block** contains the incident neutron cross section data (photon production cross sections are given in the **SIGP Block** and particle production cross sections are given in **SIGH Block**). The format of the **SIG Block** is given in Table 16. The starting index LXS of the **SIG Block** is given by $\text{JXS}(7)$. The cross section data for each reaction begins at an index defined by the corresponding relative locator from the **LSIG Block**, which are given in Table 17.

Table 16: [SIG Block](#).

Location in XSS	Description
LXS+LOCA ₁ -1	Cross section array for reaction MT ₁
LXS+LOCA ₂ -1	Cross section array for reaction MT ₂
...	
LXS+LOCA _{NMT} -1	Cross section array for reaction MT _{NMT}

Note: The number of cross section arrays NMT=NXS(4).

Table 17: Cross section array for the i -th reaction..

Location in XSS	Parameter	Description
LXS + LOCA _i -1	IE _i	Energy grid index for reaction MT _i
LXS + LOCA _i	N _{E,i}	Number of consecutive entries for MT _i
LXS + LOCA _i +1	$\sigma_i[E(l)]$ for $l = \text{IE}_i, \dots, \text{IE}_i + N_{E,i} - 1$	Cross section for reaction MT _i

Note: The energy grid, $E(l)$ is given in the [ESZ Block](#).

The energy grid index IE_i corresponds to the first energy in the grid at which a cross section is given. The MT_is are defined in the [MTR Block](#).

4.3.9 LAND, LANDP and LANDH Blocks

The [LAND Block](#) (for incident neutron reactions), [LANDP Block](#) (for photon production reactions) and [LANDH Block](#) (for particle production reactions), give the locators for the angular distribution array for each reaction MT. A locator is a *relative* index in the XSS array where some piece of data can be found. In this case, the data are the angular distributions. The format of the [LAND Block](#), [LANDP Block](#) and [LANDH Block](#) is given in Table 19 and Table 20.

The starting index LAND depends on whether it is the [LAND Block](#), [LANDP Block](#) or [LANDH Block](#) and is given in Table 18. For the particle production [LANDH Block](#), i refers to the index of the corresponding particle type defined on the [PTYPE Block](#) and is between 1 and NTYPE.

Block	LAND	NMT
LAND	JXS(8)	NXS(5)
LANDP	JXS(16)	NXS(6)
LANDH	XSS(JXS(32)+10*(i-1)+5)	XSS(JXS(31)+i-1)

Table 18: LAND and NMT values for the [TYR Block](#) and [TYRH Block](#).

The MTs are given in the [MTR Block](#), the [MTRP Block](#) and [MTRH Block](#) respectively for the [LAND Block](#), the [LANDP Block](#) and [LANDH Block](#) respectively. LOCB_i must be monotonically increasing. All locators (LOCB) are *relative* to $\text{AND}=\text{JXS}(9)$ for the [LAND Block](#), *relative* to $\text{ANDP}=\text{JXS}(17)$ for the [LANDP Block](#) and *relative* to $\text{ANDH}=\text{XSS}(\text{JXS}(32)+10*(i-1)+6)$ for the [LANDH Block](#) for particle index i .

Table 19: LAND Block.

Location in XSS	Parameter	Description
LAND	LOCB_1	Location of angular distributions for elastic scattering
LAND+1	LOCB_2	Location of angular distributions for reaction MT_1
...		
LAND+NMT	$\text{LOCB}_{\text{NMT}+1}$	Location of angular distributions for reaction MT_{NMT}

Table 20: LANDP and LANDH Block.

Location in XSS	Parameter	Description
LAND	LOCB_1	Location of angular distributions for reaction MT_1
LAND+1	LOCB_2	Location of angular distributions for reaction MT_2
...		
LAND+NMT-1	LOCB_{NMT}	Location of angular distributions for reaction MT_{NMT}

4.3.10 AND, ANDP and ANDH Blocks

The [AND Block](#), [ANDP Block](#) and [ANDH Block](#) contains angular distribution data for all reactions that produce secondary particles (neutrons for the [AND Block](#), photons for the [ANDP Block](#) and a specific particle for the [ANDH Block](#)). The format of these blocks is given in Table 22 and Table 23. The angular distribution data begins at the index specified by the locator LOCB from the [LAND Block](#), [LANDP Block](#) or [LANDH Block](#). If $\text{LOCB}_i=0$, no angular distribution data are given for reaction i and isotropic scattering is assumed in either the Lab or center-of-mass system. If $\text{LOCB}_i=-1$ no angular distribution data is given for reaction i (this can only happen for the the [AND Block](#) or [ANDH Block](#)). In this case, the angular distribution data are specified through $\text{law}=44$ in the [DLW Block](#) or [DLWH Block](#).

The starting index LAND depends on whether it is the [AND Block](#), [ANDP Block](#) or [ANDH Block](#) and are given in Table 21. For the particle production [LANDH Block](#), i refers to the index of the corresponding particle type defined on the [PTYPE Block](#) and is between 1 and NTYPE.

Block	LAND	NMT
AND	JXS(9)	NXS(5)
ANDP	JXS(17)	NXS(6)
ANDH	XSS(JXS(32)+10*(i-1)+6)	XSS(JXS(31)+i-1)

 Table 21: LAND and NMT values for the [AND Block](#) and [ANDH Block](#).

 Table 22: [AND Block](#).

Location in XSS	Description
LAND+LOCB ₁ -1	Angular distribution array for elastic scattering
LAND+LOCB ₂ -1	Angular distribution array for reaction MT ₁
...	
LAND+LOCB _{NMT+1} -1	Angular distribution array for reaction MT _{NMT}

Note: The format for the angular distribution of the i -th array is given in Table 24.

 Table 23: [ANDP Block](#) and [ANDH Block](#).

Location in XSS	Description
LAND+LOCB ₁ -1	Angular distribution array for reaction MT ₁
...	
LAND+LOCB _{NMT} -1	Angular distribution array for reaction MT _{NMT}

Note: The format for the angular distribution of the i -th array is given in Table 24.

 Table 24: Angular distribution array for the i -th reaction.

Location in XSS	Parameter	Description
LAND+LOCB ₁ -1	N_E	Number of energies at which angular distributions are tabulated.
LAND+LOCB _i	$E(l), l = 1, \dots, N_E$	Energy grid
...		
LAND+LOCB _i + N_E	$L_C(l), l = 1, \dots, N_E$	Location of tables associated with $E(l)$

The angular distribution arrays (Table 24) contains additional locators, L_C ; the sign of these locators is a flag:

- if LOCC₁=0, then distribution is isotropic and no further data is needed;

- if $\text{LOCC}_1 > 0$, then LOCC_1 points to a 32 equiprobable bin distribution (see Table 25);
- if $\text{LOCC}_1 < 0$, then LOCC_1 points to a tabulated angular distribution (see Table 26).

Table 25: Format for the 32 equiprobable bin distribution.

Location in XSS	Parameter	Description
$\text{LAND} + \text{LOCC}_1 - 1$	$P(1, K)$ $K = 1, \dots, 33$	32 equiprobable cosine bins for scattering at energy $E(l)$.

Table 26: Format for the tabulated angular distribution..

Location in XSS	Parameter	Description
$\text{LAND} + \text{LOCC}_1 - 1$	JJ	Interpolation flag [†]
$\text{LAND} + \text{LOCC}_1 $	N_P	Number of points in the distribution
$\text{LAND} + \text{LOCC}_1 + 1$	$CS_{\text{out}}(j), j = 1, \dots, N_P$	Cosine scattering angular grid
$\text{LAND} + \text{LOCC}_1 + 1 + N_P$	$\text{PDF}(j), j = 1, \dots, N_P$	Probability density function
$\text{LAND} + \text{LOCC}_1 + 1 + 2N_P$	$\text{CDF}(j), j = 1, \dots, N_P$	Cumulative density function

[†] 1 histogram interpolation,
 2 linear-linear interpolation

The [AND Block](#) and [ANDH Block](#) can use both options (either a 32 equiprobable bin or tabulated distribution). The [ANDP Block](#) on the other hand can only use 32 equiprobable bin distributions.

4.3.11 LDLW, LDLWP, DNEDL and LDLWH Blocks

The [LDLW Block](#), [LDLWP Block](#) and [LDLWH Block](#) give the locators for the energy distribution for every reaction that produces secondary neutron, secondary photons or other secondary particles (respectively). The [DNEDL Block](#) on the other hand gives the locators for the delayed neutron energy distribution for each precursor group.

The format of the [LDLW Block](#) (for secondary neutrons), the [LDLW Block](#) (for secondary photons), the [LDLWH Block](#) (for secondary particles) and the [DNEDL Block](#) (for delayed neutrons) is given in Table 28. The format for the distributions is given in Section 4.3.12.

The starting index LED depends on whether it is the [LDLW Block](#), [LDLWP Block](#), [LDLWH Block](#) or [DNEDL Block](#) and are given in Table 27. For the particle production [LDLWH Block](#), i refers to the index of the corresponding particle type defined on the [PTYPE Block](#) and is between 1 and NTYPE. These blocks are given only if the starting index, LED, is different from zero.

Block	LED	NMT
LDLW	JXS(10)	NXS(5)
LDLWP	JXS(18)	NXS(6)
LDLWH	XSS(JXS(32)+10*(i-1)+7)	XSS(JXS(31)+i-1)
DNEDL	JXS(26)	NXS(8)

Table 27: LED and NMT values for the [LDLW Block](#), the [LDLWP Block](#), the [LDLWH Block](#) and [DNEDL Block](#).

Table 28: LDLW Block.

Location in XSS	Parameter	Description
LED	LOCC ₁	Location of energy distribution data for reaction MT ₁ or group 1 (if delayed neutron)
LED+1	LOCC ₂	Location of energy distribution data for reaction MT ₂ or group 2 (if delayed neutron)
...		
LED+NMT-1	LOCC _{NMT}	Location of energy distribution data for reaction MT _{NMT} or group NMT (if delayed neutron)

Note: The LOCC_i must be monotonically increasing.

The MTs are given in the [MTR Block](#), the [MTRP Block](#) and [MTRH Block](#) respectively for the [LDLW Block](#), the [LDLWP Block](#) and [LDLWH Block](#) respectively. LOCC_i must be monotonically increasing. All locators (LOCC) are *relative* to JED=JXS(19) for the [LDLW Block](#), *relative* to JED=JXS(19) for the [LDLWP Block](#) and *relative* to JED=XSS(JXS(32)+10*(i-1)+8) for the [LDLWH Block](#) for particle index i .

4.3.12 DLW, DLWP, DLWH and DNED Blocks

The [DLW Block](#) contains secondary neutron energy distributions for all reactions producing secondary neutrons (except for elastic scattering), the [DLWP Block](#) contains secondary photon energy distributions for all photon-producing reactions, the [DLWH Block](#) contains secondary particle energy distributions for all secondary particle producing reactions and the [DNED Block](#) contains the energy distributions for the delayed neutrons. The [DLW Block](#), [DLWP Block](#), [DLWH Block](#) and [DNED Block](#) block have the same format (although there may be restrictions on which laws are allowed in these blocks). The energy distributions are given starting with a locator, LOCC, which were given in the [LDLW Block](#), [LDLWP Block](#) or [DNEDL Block](#). The locators are relative to the JED parameter. The value for JED and NMT (the number of reactions or the number of delayed precursor groups) is dependent on whether it is the [DLW Block](#), [DLWP Block](#), [DLWH Block](#) or [DNED Block](#). These values are given in Table 29. For the particle production [DLWH Block](#), i refers to the

index of the corresponding particle type defined on the [PTYPE Block](#) and is between 1 and NTYPE.

Block	JED	NMT
DLW	JXS(11)	NXS(5)
DLWP	JXS(19)	NXS(6)
DLWH	XSS(JXS(32)+10*(i-1)+8)	XSS(JXS(31)+i-1)
DNED	JXS(27)	NXS(8)

Table 29: JED and NMT for the [DLW Block](#), [DLWP Block](#) and [DLWH Block](#).

Table 30: [DLW Block](#).

Location in XSS	Description
JED+LOCC ₁ -1	Energy distribution array for reaction MT ₁ or group 1 (if delayed neutron)
JED+LOCC ₂ -1	Energy distribution array for reaction MT ₂ or group 2 (if delayed neutron)
...	
JED+LOCC _{NMT} -1	Energy distribution array for reaction MT _{NMT} or group NMT (if delayed neutron)

The i -th array has the form shown in

Table 31: Format for the secondary energy distribution..

Location in XSS	Parameter	Description
JED+LOCC _i -1	LNW ₁	Location of next law [†] relative to JED
JED+LOCC _i	LAW ₁	Name of this law
JED+LOCC _i +1	IDAT ₁	Location of data for this law relative to JED
JED+LOCC _i +2	N_R	Number of interpolation regions to define law applicability regime
JED+LOCC _i +3	$\text{NBT}(l), l = 1, \dots, N_R$	ENDF interpolation parameters
JED+LOCC _i +3+ N_R	$\text{INT}(l), l = 1, \dots, N_R$	ENDF interpolation scheme [‡]
JED+LOCC _i +3+2 N_R	N_E	Number of energies
JED+LOCC _i +4+2 N_R	$E(l), l = 1, \dots, N_E$	Tabulated energy points
JED+LOCC _i +4+2 N_R + N_E	$P(l), l = 1, \dots, N_E$	Probability of law validity [*]
JED+IDAT ₁ - 1	LDAT(l), $l = 1, \dots, L$	Law data for LAW ₁ .
JED+LNW ₁ - 1	LNW ₂	Location of next law relative to JED

Continued on next page

Table 31: Format for the secondary energy distribution. (continued)

Location in XSS	Parameter	Description
JED+LNW ₁	LAW ₂	Name of this law
JED+LNW+1	IDAT ₂	Location of data for this law relative to JED
...		

[†] If LNW_{*i*} = 0 then LAW₁ is used regardless of other circumstances.

[‡] If $N_R = 0$, NBT and INT are omitted and linear-linear interpolation is assumed.

* If the particle energy $E < E(1)$, then $P(E) = P(1)$. If $E > E(N_E)$, then $P(E) = P(N_E)$. If more than one law is given, then LAW₁ is used only if $\xi < P(E)$ where ξ is a random number between 0 and 1.

The format for the law data depends on the law. The length, L , of the law data array, LDAT, is determined from parameters with LDAT. The various LDAT arrays and their formats are given in the following tables. Laws 2 (Table 33) and 4 (Table 35) are used to describe spectra of secondary photons from neutron collisions. All laws—except for Law 2—are used to describe the spectra of scattered neutrons.

In the following tables, we provide relative locations of data in the LDAT array rather than the absolute locations in the XSS array. Table 31 defines the starting location of the LDAT array within the XSS array.

TODO: add law=33 and other charged particle related laws, specify which laws are used for which type of outgoing particle

4.3.12.1 LAW=1—Tabular Equiprobable Energy Bins

Table 32: LAW=1 (From ENDF Law 1).

Location	Parameter	Description
LDAT(1)	N_R	Number of interpolation regions between tables of E_{out}
LDAT(2)	$\text{NBT}(l), l = 1, \dots, N_R$	ENDF interpolation parameters
LDAT(2+ N_R)	$\text{INT}(l), l = 1, \dots, N_R$	ENDF interpolation scheme [†]
LDAT(2+2 N_R)	N_E	Number of incident energies tabulated
LDAT(3+2 N_R)	$E_{\text{in}}(l), l = 1, \dots, N_E$	List of incident energies for which E_{out} is tabulated
LDAT(3+2 $N_R + N_E$)	NET	Number of outgoing energies in each E_{out} table
LDAT(4+2 * $N_R + N_E$)	$E_{\text{out}_1}(l), l = 1, \dots, \text{NET}$ $E_{\text{out}_2}(l), l = 1, \dots, \text{NET}$...	E_{out} tables [‡]
	$E_{\text{out}_{N_E}}(l), l = 1, \dots, \text{NET}$	

Continued on next page

Table 32: LAW=1 (From ENDF Law 1) (continued)

Location	Parameter	Description
----------	-----------	-------------

[†] If $N_R = 0$, NBT and INT are omitted and linear-linear interpolation is assumed.

[‡] E_{out} tables consist of NET boundaries of NET-1 equally likely energy intervals. Linear-linear interpolation is used between intervals.

4.3.12.2 LAW=2—Discrete Photon Energy

Table 33: LAW=2—Discrete Photon Energy.

Location	Parameter	Description
LDAT(1)	LP	Indicator of whether the photon is a primary or non-primary photon
LDAT(2)	EG	Photon energy or binding energy

Note: If LP=0 or LP=1, the photon energy is EG. If LP=2, the photon energy is

$$\text{EG} + \left(\frac{\text{AWR}}{\text{AWR} + 1} \right) E_N$$

where AWR is the atomic weight ratio and E_N is the incident neutron energy.

4.3.12.3 LAW=3—Level Scattering

Table 34: LAW=3—Level Scattering.

Location	Parameter	Description
LDAT(1)		$(A + 1)/A Q $
LDAT(2)		$(A/(A + 1))^2$

The outgoing center-of-mass energy is calculated as:

$$E_{\text{out}}^{\text{CM}} = \text{LDAT}(2) * (E - \text{LDAT}(1)), \quad (2)$$

where

$E_{\text{out}}^{\text{CM}}$ = outgoing center-of-mass energy

E = incident energy

A = atomic weight ratio

Q = Q -value

The outgoing neutron energy in the laboratory system is:

$$E_{\text{out}}^{\text{LAB}} = E_{\text{out}}^{\text{CM}} + \left\{ E + 2\mu_{\text{CM}}(A+1)(E \cdot E_{\text{out}}^{\text{CM}})^{1/2} \right\} / (A+1)^2 \quad (3)$$

where μ_{CM} is the cosine of the center-of-mass scattering angle

4.3.12.4 LAW=4—Continuous Tabular Distribution

Table 35: LAW=4 (From ENDF-6 LAW=1).

Location	Parameter	Description
LDAT(1)	N_R	The number of interpolation regions
LDAT(2)	$\text{NBT}(l), l = 1, \dots, N_R$	ENDF interpolation parameters
LDAT(2+ N_R)	$\text{INT}(l), l = 1, \dots, N_R$	ENDF interpolation scheme [†]
LDAT(2+2 N_R)	N_E	Number of energies at which distributions are tabulated
LDAT(3+2 N_R)	$E(l), l = 1, \dots, N_E$	Incident neutron energies
LDAT(3+2 $N_R + N_E$)	$\text{L}(l), l = 1, \dots, N_E$	Locations of distributions [‡]

[†] If $N_R = 0$, NBT and INT are omitted and linear-linear interpolation is assumed.

[‡] Relative to JXS(11) (neutron reactions), JXS(19) (photon-producing reactions), or JXS(27) (delayed neutrons).

The data associated with each incident neutron energy begins at the location $\text{L}(l)$. The format for the data is given in Table 36, where for $E(1)$ let $\text{K}=3+2N_R + 2N_E$.

Table 36: Secondary energy distribution for each incident energy in LAW=4..

Location	Parameter	Description
Data for E(1)		
LDAT(K)	INTT'	Interpolation parameter
LDAT(K+1)	N_p	Number of points in the distribution
LDAT(K+2)	$E_{\text{out}}(l), l = 1, \dots, N_p$	Outgoing energy grid
LDAT(K+2 + N_p)	$\text{PDF}(l), l = 1, \dots, N_p$	Probability Density Function
LDAT(K+2 + 2 N_p)	$\text{CDF}(l), l = 1, \dots, N_p$	Cumulative Density Function
Data for E(2)—same format for E(1)		
...		
Data for E(N_E)—same format for E(1)		

Combined interpolation parameter. The first element in the data is INTT' or the interpolation parameter, which is a combination of two other parameters:

1. the number of discrete photon lines, N_D , and
2. the interpolation scheme for the subsequent data, INTT , which has two valid values:
 $\text{INTT}=1$ histogram distribution, and
 $\text{INTT}=2$ linear-linear distribution.

Given the definition of N_D and INTT , the interpolation parameter, INTT' , is defined as the combination of N_D and INTT :

$$\text{INTT}' = 10N_D + \text{INTT}. \quad (4)$$

Since N_D describe the number of *discrete* photon lines, the remaining $(N_p - N_D)$ values describe a continuous distribution. In this way, the distribution may be discrete, continuous, or a discrete distribution superimposed upon a continuous background.

4.3.12.5 LAW=5—General Evaporation Spectrum

Table 37: LAW=5 (From ENDF-6, MF=5, LF=5).

Location	Parameter	Description
LDAT(1)	N_R	Interpolation scheme between T 's
LDAT(2)	$\text{NBT}(l), l = 1, \dots, N_R$	
LDAT(2+ N_R)	$\text{INT}(l), l = 1, \dots, N_R$	
LDAT(2+2 N_R)	N_E	Number of incident energies tabulated
LDAT(3+2 N_R)	$E(l), l = 1, \dots, N_E$	Incident energy table
LDAT(3+2 $N_R + N_E$)	$\theta(l), l = 1, \dots, N_E$	Effective temperature tabulated on incident energies
LDAT(3+2 $N_R + 2N_E$)	NET	Number of X 's tabulated
LDAT(4+2 $N_R + 2N_E$)	$X(l), l = 1, \dots, \text{NET}$	Equiprobable bins

$$E_{\text{out}} = X(\xi)\theta(E) \quad (5)$$

where:

- $X(\xi)$ is a randomly sampled table of X 's;
- $\theta(E)$ is the effective temperature tabulated on incident energy; and
- E is the incident energy.

4.3.12.6 LAW=7—Simple Maxwellian Fission Spectrum

Table 38: LAW=7 (From ENDF-6, MF=5, LF=7).

Location	Parameter	Description
LDAT(1)	N_R	Interpolation scheme between T 's
LDAT(2)	$\text{NBT}(l), l = 1, \dots, N_R$	
LDAT(2+ N_R)	$\text{INT}(l), l = 1, \dots, N_R$	
LDAT(2+2 N_R)	N_E	Number of incident energies tabulated
LDAT(3+2 N_R)	$E(l), l = 1, \dots, N_E$	Incident energy table
LDAT(3+2 $N_R + N_E$)	$\theta(l), l = 1, \dots, N_E$	Effective temperature tabulated on incident energies
LDAT(3+2 $N_R + 2N_E$)	U	Restriction energy

The outgoing energy, E_{out} , can be calculated as

$$f(E \rightarrow E_{\text{out}}) = \frac{\sqrt{E_{\text{out}}}}{I} e^{-E_{\text{out}}/\theta(E)} \quad (6)$$

where:

I is the normalization constant

$$I = \theta^{3/2} \frac{\sqrt{\pi}}{2} \text{erf} \left(\sqrt{(E - U)/\theta} \right) - \sqrt{(E - U)/\theta} e^{-(E - U)/\theta}, \quad (7)$$

θ is tabulated as a function of incident energy, E ; and

U is a constant introduced to define the proper upper limit for the final particle energy such that $0 \leq E_{\text{out}} \leq (E - U)$.

4.3.12.7 LAW=9—Evaporation Spectrum

Table 39: LAW=9 (From ENDF-6, MF=5, LF=9).

Location	Parameter	Description
LDAT(1)	N_R	Interpolation scheme between T 's
LDAT(2)	$\text{NBT}(l), l = 1, \dots, N_R$	
LDAT(2+ N_R)	$\text{INT}(l), l = 1, \dots, N_R$	
LDAT(2+2 N_R)	N_E	Number of incident energies tabulated
LDAT(3+2 N_R)	$E(l), l = 1, \dots, N_E$	Incident energy table
LDAT(3+2 $N_R + N_E$)	$\theta(l), l = 1, \dots, N_E$	Effective temperature tabulated on incident energies

Continued on next page

Table 39: LAW=9 (From ENDF-6, MF=5, LF=9) (continued)

Location	Parameter	Description
LDAT(3+2N _R + 2N _E)	U	Restriction energy

The outgoing energy, E_{out} , can be calculated as

$$f(E \rightarrow E_{\text{out}}) = \frac{\sqrt{E_{\text{out}}}}{I} e^{-E_{\text{out}}/\theta(E)} \quad (8)$$

where:

I is the normalization constant

$$I = \theta^2 \left[1 - e^{-(E-U)/\theta} \left(1 + \frac{E-U}{\theta} \right) \right], \quad (9)$$

θ is tabulated as a function of incident energy, E ; and

U is a constant introduced to define the proper upper limit for the final particle energy such that

$$0 \leq E_{\text{out}} \leq (E - U).$$

Note: Equation 8 is the same as Equation 6; just the definitions of I in Equation 7 and Equation 9 are different.

4.3.12.8 LAW=11—Energy Dependent Watt Spectrum

Table 40: LAW=11 (From ENDF-6, MF=5, LF=11).

Location	Parameter	Description
LDAT(1)	N_{R_a}	Interpolation scheme between a 's
LDAT(2)	$\text{NBT}_a(l), l = 1, \dots, N_{R_a}$	
LDAT(2+N _{R_a})	$\text{INT}_a(l), l = 1, \dots, N_{R_a}$	
LDAT(2+2N _{R_a})	N_{E_a}	Number of incident energies tabulated for $a(E_{\text{in}})$ table
LDAT(3+2N _{R_a})	$E_a(l), l = 1, \dots, N_{E_a}$	Incident energy table
LDAT(3+2N _{R_a} + N _{E_a})	$a(l), l = 1, \dots, N_{E_a}$	Tabulated a 's
let L = 3 + 2(N _{R_a} + N _{E_a})		
LDAT(L)	N_{R_b}	Interpolation scheme between b 's
LDAT(L+1)	$\text{NBT}_b(l), l = 1, \dots, N_{R_b}$	
LDAT(L+1+N _{R_b})	$\text{INT}_b(l), l = 1, \dots, N_{R_b}$	
LDAT(L+1+2N _{R_b})	N_{E_b}	Number of incident energies tabulated for $b(E_{\text{in}})$ table

Continued on next page

Table 40: LAW=11 (From ENDF-6, MF=5, LF=11) (continued)

Location	Parameter	Description
L DAT(L+2+2N _{R_b})	$E_b(l), l = 1, \dots, N_{E_b}$	Incident energy table
L DAT(L+2+2N _{R_b} + N _{E_b})	$b(l), l = 1, \dots, N_{E_b}$	Tabulated b 's
L DAT(L+2+2N _{R_b} + 2N _{E_b})	U	Rejection energy

The outgoing energy, E_{out} , can be calculated as

$$f(E \rightarrow E_{\text{out}}) = \frac{e^{-E_{\text{out}}/a}}{I} \sinh\left(\sqrt{bE_{\text{out}}}\right) \quad (10)$$

where:

I is the normalization constant

$$I = \frac{1}{2} \sqrt{\frac{\pi a^3 b}{4}} e^{(ab/4)} \left[\operatorname{erf}\left(\sqrt{\frac{E-U}{a}} - \sqrt{\frac{ab}{4}}\right) + \operatorname{erf}\left(\sqrt{\frac{E-U}{a}} + \sqrt{\frac{ab}{4}}\right) \right] - ae^{-(E-U)/a} \sinh \sqrt{b(E-U)}; \quad (11)$$

a and b are tabulated energy-dependent parameters; and

U is a constant introduced to define the proper upper limit for the final particle energy such that $0 \leq E_{\text{out}} \leq (E - U)$.

4.3.12.9 LAW=22—Tabular Linear Functions of Incident Energy Out

Table 41: LAW=22 (From UK Law 2).

Location	Parameter	Description
L DAT(1)	N_R	Interpolation parameters
L DAT(2)	$\text{NBT}(l), l = 1, \dots, N_R$	
L DAT(2+N _R)	$\text{INT}(l), l = 1, \dots, N_R$	
L DAT(2+2N _R)	N_E	Number of incident energies tabulated
L DAT(3+2N _R)	$E_{\text{in}}(l), l = 1, \dots, N_E$	Tabulated incident energies for E_{out} tables
L DAT(3+2N _R + N _E)	$\text{LOCE}(l), l = 1, \dots, N_E$	Locators of E_{out} tables
Data for $E_{\text{in}}(1)$ Let $L = 3 + 2N_R + 2N_E$:		
L DAT(L)	NF_1	
L DAT(L+1)	$P_{1k}, k = 1, \dots, \text{NF}_1$	
L DAT(L+1+NF ₁)	$T_{1k}, k = 1, \dots, \text{NF}_1$	

Continued on next page

Table 41: LAW=22 (From UK Law 2) (continued)

Location	Parameter	Description
LDAT(L+1+2NF ₁)	$C_{1k}, k = 1, \dots, \text{NF}_1$	
Data for $E_{\text{in}}(2)$:		
...		

Tables of P_{ik} , C_{ik} , and T_{ik} are given at a number of incident energies, E_{in} . If

$$E_{\text{in}}(l) \leq E < E_{\text{in}}(l+1) \quad (12)$$

then the secondary neutron energy is:

$$E_{\text{out}} = C_{ik}(E - T_{ik}), \quad (13)$$

where k is chosen according to

$$\sum_{j=1}^k P_{ij} < \xi \leq \sum_{k=1}^{k+1} P_{ij} \quad (14)$$

for a given random number, $\xi \in [0, 1)$.

4.3.12.10 LAW=24—Tabular Energy Multipliers

Table 42: LAW=24 (From UK Law 6).

Location	Parameter	Description
LDAT(1)	N_R	Interpolation scheme between T 's
LDAT(2)	$\text{NBT}(l), l = 1, \dots, N_R$	
LDAT(2+ N_R)	$\text{INT}(l), l = 1, \dots, N_R$	
LDAT(2+2 N_R)	N_E	Number of incident energies tabulated
LDAT(3+2 N_R)	$E_{\text{in}}(l), l = 1, \dots, N_E$	List of incident energies for which T is tabulated
LDAT(3+2 N_R + N_E)	NET	Number of outgoing values in each table
LDAT(4+2 N_R + N_E)	$T_1(l), l = 1, \dots, \text{NET}$	Tables have NET boundaries with $\text{NET}-1$ equally likely intervals. Linear-linear interpolation is used between intervals.
	$T_2(l), l = 1, \dots, \text{NET}$	
	...	
	$T_{N_E}(l), l = 1, \dots, \text{NET}$	

The outgoing energy, E_{out} can be calculated as:

$$E_{\text{out}} = T_k(l) * E \quad (15)$$

where:

$T_k(l)$ is sampled from the tables and
 E is the incident energy.

4.3.12.11 LAW=44—Kalbach-87 Formalism

Table 43: LAW=44 (From ENDF-6 MF=6 LAW=1, LANG=2).

Location	Parameter	Description
LDAT(1)	N_R	Interpolation scheme between tables of E_{out}
LDAT(2)	$\text{NBT}(l), l = 1, \dots, N_R$	ENDF interpolation parameters
LDAT(2+ N_R)	$\text{INT}(l), l = 1, \dots, N_R$	ENDF interpolation scheme [†]
LDAT(2+2 N_R)	N_E	Number of energies at which distributions are tabulated
LDAT(3+2 N_R)	$E(l), l = 1, \dots, N_E$	Incident neutron energies
LDAT(3+2 $N_R + N_E$)	$L(l), l = 1, \dots, N_E$	Locations of distributions [‡]

[†] If $N_R = 0$, NBT and INT are omitted and linear-linear interpolation is assumed.

[‡] Relative to JXS(11) (neutron reactions), JXS(19) (photon-producing reactions), or JXS(27) (delayed neutrons).

The data associated with each incident neutron energy begins at the location $L(l)$. The format for the data is given in Table 44, where for $E(1)$ let $K=3+2N_R + 2N_E$.

Table 44: Secondary energy distribution for each incident energy in LAW=44.

Location	Parameter	Description
Data for E(1)		
LDAT(K)	INTT'	Interpolation parameter
LDAT(K+1)	N_p	Number of points in the distribution
LDAT(K+2)	$E_{\text{out}}(l), l = 1, \dots, N_p$	Outgoing energy grid
LDAT(K+2 + N_p)	$\text{PDF}(l), l = 1, \dots, N_p$	Probability Density Function
LDAT(K+2 + 2 N_p)	$\text{CDF}(l), l = 1, \dots, N_p$	Cumulative Density Function
LDAT(K+2 + 3 N_p)	$R(l), l = 1, \dots, N_p$	Precompound fraction r
LDAT(K+2 + 4 N_p)	$A(l), l = 1, \dots, N_p$	Angular distribution slope value a
Data for E(2)—same format for E(1)		
...		
Data for E(N_E)—same format for E(1)		

The interpolation parameter, INTT' has the same definition as in LAW=4, described in Paragraph 4.3.12.4.

The angular distributions for neutrons are sampled from:

$$p(\mu, E_{\text{in}}, E_{\text{out}}) = \frac{1}{2} \frac{a}{\sinh(a)} [\cosh(a\mu) + r \sinh(a\mu)]. \quad (16)$$

4.3.12.12 LAW=61—Like LAW=44, but tabular angular distribution instead of Kalbach-87

Table 45: LAW=61.

Location	Parameter	Description
LDAT(1)	N_R	Number of interpolation regions
LDAT(2)	$\text{NBT}(l), l = 1, \dots, N_R$	ENDF interpolation parameters
LDAT(2+ N_R)	$\text{INT}(l), l = 1, \dots, N_R$	ENDF interpolation scheme [†]
LDAT(2+2 N_R)	N_E	Number of energies at which distributions are tabulated
LDAT(3+2 N_R)	$E(l), l = 1, \dots, N_E$	Incident neutron energies
LDAT(3+2 $N_R + N_E$)	$L(l), l = 1, \dots, N_E$	Locations of distributions [‡]

[†] If $N_R = 0$, NBT and INT are omitted and linear-linear interpolation is assumed.

[‡] Relative to JXS(11) (neutron reactions), JXS(19) (photon-producing reactions), or JXS(27) (delayed neutrons).

The data associated with each incident neutron energy begins at the location $L(l)$. The format for the data is given in Table 46, where for $E(1)$ let $K=3+2N_R + 2N_E$.

Table 46: Secondary energy distribution for each incident energy in LAW=61.

Location	Parameter	Description
Data for E(1)		
LDAT(K)	INTT'	Interpolation parameter
LDAT(K+1)	N_p	Number of points in the distribution
LDAT(K+2)	$E_{\text{out}}(l), l = 1, \dots, N_p$	Outgoing energy grid
LDAT(K+2 + N_p)	$\text{PDF}(l), l = 1, \dots, N_p$	Probability Density Function
LDAT(K+2 + 2 N_p)	$\text{CDF}(l), l = 1, \dots, N_p$	Cumulative Density Function
LDAT(K+2 + 3 N_p)	$\text{LC}(l), l = 1, \dots, N_p$	Location of tables associated with incident energies $E(l)$. See Table 47
Data for E(2)—same format for E(1)		
...		
Data for E(N_E)—same format for E(1)		

The interpolation parameter, INTT' has the same definition as in LAW=4, described in Paragraph 4.3.12.4.

The J -th array for the tabular angular distribution has the form shown in Table 47. For the angular distribution, the locators L are relative to $JXS(11)$ for neutron reactions or $JXS(19)$ for photon-producing reactions. Thus,

$$\begin{aligned} L &= JXS(11) + |LC(J)| - 1 \text{ (for neutron reactions),} \\ L &= JXS(19) + |LC(J)| - 1 \text{ (for photon-producing reactions).} \end{aligned}$$

Table 47: Angular distribution for LAW=61.

Location	Parameter	Description
LDAT(L+1)	JJ	Interpolation flag
LDAT(L+2)	N_P	Number of points in the distribution
LDAT(L+3)	$CS_{\text{out}}(j), j = 1, \dots, N_P$	Cosine scattering angular grid
LDAT(L+3+ N_P)	$PDF(j), j = 1, \dots, N_P$	Probability density function
LDAT(L+3+2 N_P)	$CDF(j), j = 1, \dots, N_P$	Cumulative density function

Valid values for the interpolation flag, JJ, are the same as for INTT:

JJ=1 histogram distribution, and

JJ=2 linear-linear distribution.

4.3.12.13 LAW=66— N -body phase space distribution

Table 48: LAW=66 (From ENDF-6 MF=6 LAW=6).

Location	Parameter	Description
LDAT(1)	NPSX	Number of bodies in the phase space
LDAT(2)	A_P	Total mass ratio for the NPSX particles.
LDAT(3)	INTT	Interpolation parameter [†]
LDAT(4)	N_P	Number of points in the distribution
LDAT(5)	$\xi_{\text{out}}(j), j = 1, \dots, N_P$	ξ grid (between 0 and 1)
LDAT(5+ N_P)	$PDF(j), j = 1, \dots, N_P$	Probability density function
LDAT(5+2 N_P)	$CDF(j), j = 1, \dots, N_P$	Cumulative density function

[†] **INTT=1** histogram distribution,
INTT=2 linear-linear distribution.

The outgoing energy is

$$E_{\text{out}} = T(\xi)E_i^{\text{max}} \quad (17)$$

where

$$E_i^{\max} = \frac{A_p - 1}{A_p} \left(\frac{A}{A + 1} E_{\text{in}} + Q \right) \quad (18)$$

and $T(\xi)$ is sampled from:

$$P_i(\mu, E_{\text{in}}, T) = C_n \sqrt{T} (E_i^{\max} - T)^{3n/2-4} \quad (19)$$

While MCNP will only use the values for NPSX and A_P (given in LDAT(1) and LDAT(2)), NJOY does add the derived probability density function and cumulative density function for $T(\xi)$ starting at LDAT(3).

4.3.12.14 LAW=67—Laboratory Angle-Energy Law

Table 49: LAW=67 (From ENDF-6 MF=6 LAW=7).

Location	Parameter	Description
LDAT(1)	N_R	Number of interpolation regions
LDAT(2)	$\text{NBT}(l), l = 1, \dots, N_R$	ENDF interpolation parameters
LDAT(2+ N_R)	$\text{INT}(l), l = 1, \dots, N_R$	ENDF interpolation scheme [†]
LDAT(2+2 N_R)	N_E	Number of energies at which distributions are tabulated
LDAT(3+2 N_R)	$E(l), l = 1, \dots, N_E$	Incident neutron energies
LDAT(3+2 $N_R + N_E$)	$L(l), l = 1, \dots, N_E$	Locations of distributions [‡]

[†] If $N_R = 0$, NBT and INT are omitted and linear-linear interpolation is assumed.

[‡] Relative to JXS(11) (neutron reactions), JXS(19) (photon-producing reactions), or JXS(27) (delayed neutrons).

The data associated with each distribution begins at location $L(l)$. The format for the data is given in Table 50, where for $E(1)$ let $K = 3 + 2N_R + 2N_E$.

Table 50: Angular distribution for LAW=67.

Location	Parameter	Description
LDAT(K)	INTMU	Interpolation scheme [†]
LDAT(K+1)	NMU	Number of secondary cosines
LDAT(K+2)	$\text{XMU}(l), l = 1, \dots, \text{NMU}$	Secondary cosines

Continued on next page

Table 50: Angular distribution for LAW=67 (continued)

Location	Parameter	Description
L DAT(K+2+N MU)	L MU(l), $l = 1, \dots, \text{N MU}$	Locations of data for each secondary cosine. See Table 51

† **INTMU=1** histogram distribution,
INTMU=2 linear-linear distribution.

The format for the secondary energy distribution (for each cosine bin, X MU) is given in Table 51. For the energy distribution, the locators, L MU, are relative to J XS(11) or J XS(19). Thus,

$$\begin{aligned} L_l &= \text{J XS}(11) + \text{L MU}(l) \text{ (for neutron reactions),} \\ L_l &= \text{J XS}(19) + \text{L MU}(l) \text{ (for photon-producing reactions).} \end{aligned}$$

Table 51: Secondary energy distribution for each cosine bin in LAW=67.

Location	Parameter	Description
L DAT(L_l)	INTEP	Interpolation parameter between secondary energies†
L DAT($L_l + 1$)	NPEP	Number of secondary energies
L DAT($L_l + 2$)	$E_P(l)$, $l = 1, \dots, \text{NPEP}$	Secondary energy grid
L DAT($L_l + 2 + \text{NPEP}$)	PDF(l), $l = 1, \dots, \text{NPEP}$	Probability density function
L DAT($L_l + 2 + 2\text{NPEP}$)	CDF(l), $l = 1, \dots, \text{NPEP}$	Cumulative density function

† **INTEP=1** histogram distribution,
INTEP=2 linear-linear distribution.

4.3.12.15 Energy-Dependent Neutron Yields

There are additional numbers to be found for neutrons in the DLW Block and DLWP Block. For those reactions with entries in the TYR Block that are greater than 100 in absolute value, there must be neutron yields, $Y(E)$ provided as a function of neutron energy. The neutron yields are handled similarly to the average number of neutrons per fission, $\nu(E)$ that is given for the fission reactions. These yields are a part of the coupled energy-angle distributions given in File 6 of ENDF-6 data.

The i -th array has the form given in Table 52, where $\text{KY} = \text{JED} + |\text{TY}_i| - 101$.

Table 52: Energy-Dependent Neutron Yields.

Location	Parameter	Description
KY	N_R	Number of interpolation regions
KY+1	$\text{NBT}(l), l = 1, \dots, N_R$	ENDF interpolation parameters
KY+1+ N_R	$\text{INT}(l), l = 1, \dots, N_R$	ENDF interpolation scheme [†]
KY+1+2 N_R	N_E	Number of energies
KY+2+2 N_R	$E(l), l = 1, \dots, N_E$	Tabular energy points
KY+2+ $N_R + N_E$	$Y(l), l = 1, \dots, N_E$	Corresponding energy-dependent yields

[†] If $N_R = 0$, NBT and INT are omitted and linear-linear interpolation is assumed.

4.3.13 GPD Block

The **GPD Block** contains the *total* photon production cross section, tabulated on the energy grid given in the **ESZ Block**, the size of which is given by `NXS(3)`. The **GPD Block** only exists if `JXS(12) ≠ 0` and is shown in Table 53.

Table 53: GPD Block.

Location in XSS	Parameter	Description
S_{GPD}	$\sigma_\gamma(l), l = 1, \dots, \text{NES}$	Total photon production cross section

In addition to the total photon production cross section, the outgoing photon energies *may* be given.⁴ There are 30 groups for the incident neutron energies, the boundaries of which are shown in Table 54.

For each incident neutron energy group, the outgoing photon energies are discretized into 20 equiprobable energy groups, thus creating a 30×20 matrix. The outgoing energies are given in the **GPD Block**, after the total photon production cross section as shown in Table 55. Note that this matrix is only used for older tables that do not provide expanded photon production data. The format of this Block is given in Table 55. The XSS array index at the start of the **GPD Block**, $S_{\text{GPD}} = \text{JXS}(12)$.

 Table 55: Outgoing photon energies in **GPD Block**..

Location in XSS	Parameter	Description
$S_{\text{GPD}} + \text{NES}$	$E_1(K), K = 1, 20$	20 equiprobable outgoing photon energies for incident neutron $E < E_N(2)$

Continued on next page

⁴Note that this is an obsolete format. It only exists when `JXS(12) ≠ 0` and `JXS(13) = 0`.

Table 55: Outgoing photon energies in [GPD Block](#). (continued)

Location in XSS	Parameter	Description
$S_{\text{GPD}+\text{NES}+20}$	$E_2(K), K = 1, 20$	20 equiprobable outgoing photon energies for incident neutron $E_N(2) \leq E < E_N(3)$
...		
$S_{\text{GPD}+\text{NES}+(i-1)*20}$	$E_i(K), K = 1, 20$	20 equiprobable outgoing photon energies for incident neutron $E_N(i) \leq E < E_N(i+1)$
...		
$S_{\text{GPD}+\text{NES}+(30-1)*20}$	$E_N(K), K = 1, 20$	20 equiprobable outgoing photon energies for incident neutron $E \geq E_N(30)$

4.3.14 SIGP and SIGH Blocks

The [SIGP Block](#) contains the photon production cross section data and the [SIGH Block](#) contains the particle production cross section data. The format of the [SIGP Block](#) and [SIGH Block](#) is given in Table 57. The starting index depends on whether it is the [LSIGP Block](#) or [LSIGH Block](#) and are given in Table 56. For the particle production [SIGH Block](#), i refers to the index of the corresponding particle type defined on the [PTYPE Block](#) and is between 1 and NTYPE.

The cross section data begins at the index specified by the locator, LOCA_i , given in the [LSIGP Block](#) or [LSIGH Block](#). The MTs are defined in the [MTRP Block](#) or [MTRH Block](#). All indices to the XSS array are *relative* to SIG.

Table 57: SIGP Block.

Location in XSS	Parameter	Description
$\text{SIG}+\text{LOCA}_1-1$	MFTYPE_1	Cross section array for reaction MT_1
$\text{SIG}+\text{LOCA}_2-1$	MFTYPE_2	Cross section array for reaction MT_2
...		
$\text{SIG}+\text{LOCA}_{\text{NMT}}-1$	$\text{MFTYPE}_{\text{NMT}}$	Cross section array for reaction MT_{NMT}

The format of the i -th cross section array has two possible forms depending on the first number in the array, MFTYPE.

1. If $\text{MFTYPE}=12$ or $\text{MFTYPE}=16$, yield data taken from ENDF File 12 or 6, respectively (see Table 58). With this format, the photon or particle production cross section can be constructed using Equation 20:

$$\sigma_i(E) = Y(E) * \sigma_{\text{MTMULT}}(E). \quad (20)$$

2. If $\text{MFTYPE}=13$, photon production cross section data from ENDF File 13 (see Table 59). This form is only allowed for the [SIGP Block](#).

Table 54: Discrete neutron energy boundaries.

Group #	Upper Boundary (MeV)	Group #	Upper Boundary (MeV)
1	1.39×10^{-10}	16	0.184
2	1.52×10^{-7}	17	0.303
3	4.14×10^{-7}	18	0.500
4	1.13×10^{-6}	19	0.823
5	3.06×10^{-6}	20	1.353
6	8.32×10^{-6}	21	1.738
7	2.26×10^{-5}	22	2.232
8	6.14×10^{-5}	23	2.865
9	1.67×10^{-4}	24	3.68
10	4.54×10^{-4}	25	6.07
11	1.235×10^{-3}	26	7.79
12	3.35×10^{-3}	27	10
13	9.23×10^{-3}	28	12
14	2.48×10^{-2}	29	13.5
15	6.76×10^{-2}	30	15

Block	SIG	NMT
SIGP	JXS(15)	NXS(6)
SIGH	XSS(JXS(32)+10*(i-1)+4)	XSS(JXS(31)+i-1)

 Table 56: SIG and NMT values for the [SIGP Block](#) and [SIGH Block](#).

Table 58: Photon production array if MFTYPE=12 or 16.

Location in XSS	Parameter	Description
JXS(15)+LOCA _i -1	MFTYPE	12 or 16
JXS(15)+LOCA _i	MTMULT	Neutron MT whose cross section should multiply the yield
JXS(15)+LOCA _i +1	N_R	Number of interpolation regions
JXS(15)+LOCA _i +2	$\text{NBT}(l), l = 1, \dots, N_R$	ENDF interpolation parameters [†]
JXS(15)+LOCA _i +2 + N_R	$\text{INT}(l), l = 1, \dots, N_R$	ENDF interpolation scheme
JXS(15)+LOCA _i +2 + $2 * N_R$	N_E	Number of energies at which the yield is tabulated

Continued on next page

Table 58: Photon production array if MFTYPE=12 or 16 (continued)

Location in XSS	Parameter	Description
JXS(15)+LOCA _i +3 +2 * N _R	$E(l), l = 1, \dots, N_E$	Energies
JXS(15)+LOCA _i +3 +2 * N _R + N _E	$Y(l), l = 1, \dots, N_E$	Yields

[†] If $N_R = 0$, NBT and INT are omitted and linear-linear interpolation is used.

Table 59: Photon production cross section array if MFTYPE=13.

Location in XSS	Parameter	Description
JXS(15)+LOCA _i -1	MFTYPE	13
JXS(15)+LOCA _i	IE	Energy grid index
JXS(15)+LOCA _i +1	N _E	Number of consecutive entries
JXS(15)+LOCA _i +2	$\sigma_{\gamma,i}[E(K)],$ $K = IE, \dots, IE + N_E - 1$	Photon production cross sections for reaction MT _i

Note: The MT_is are defined in the [MTRP Block](#).

4.3.15 YP and YH Blocks

The [YP Block](#) and [YH Block](#) contains a list of MT identifiers of cross sections that are used as yield multipliers in Equation 20 to calculate the photon production cross sections (for the [YP Block](#)) and the secondary particle production cross sections (for the [YH Block](#)) and are referenced by the MTMULT parameter in Table 58. The format of the [YP Block](#) and [YH Block](#) is given in Table 60.

The starting index LY depends on whether it is the [YP Block](#) or [YH Block](#). For the [YP Block](#), LY = NXS(6). For the particle production [YH Block](#), JED = XSS(JXS(32)+10*(i-1)+8) in which i refers to the index of the corresponding particle type defined on the [PTYPE Block](#) and is between 1 and NTYPE. These blocks are given only if the starting index LY is different from zero.

Table 60: YP Block.

Location in XSS	Parameter	Description
LY	NYP	Number of MTs to follow
LY+1	MTY(l), l = 1, ..., NYP	MTs.

4.3.16 FIS Block

The **FIS Block** contains the total fission cross section. The **FIS Block** exists if $\text{JXS}(21) \neq 0$, but is generally not provided; the total fission cross section is redundant as the total fission cross section is the summation of first-, second-, third-, and fourth-chance fission ($\text{MT}=19, 20, 21$, and 38);

$$\sigma_{f,t}(E) = \sigma_{(n,f)} + \sigma_{(n,nf)} + \sigma_{(n,2nf)} + \sigma_{(n,3nf)}. \quad (21)$$

The format of the **FIS Block** is given in Table 61.

Table 61: FIS Block.

Location in XSS	Parameter	Description
$\text{JXS}(21)$	IE	Energy grid index
$\text{JXS}(21)+1$	N_E	Number of consecutive entries
$\text{JXS}(21)+2$	$\sigma_f[E(l)], K = \text{IE}, \dots, \text{IE} + N_E - 1$	Total fission cross sections

Note: The energy $E(l)$ is given in the **ESZ Block**.

4.3.17 UNR Block

The **UNR Block** contains the unresolved resonance range probability tables. It exists if $\text{JXS}(23) \neq 0$ and begins at location $\text{JXS}(23)$ in XSS. The **UNR Block** has several flags that have special meaning:

ILF The ILF flag is the inelastic competition flag.

ILF < 0 The inelastic cross section is zero within the entire unresolved energy range.

ILF > 0 The value of ILF is a special MT number whose tabulation is the sum of the inelastic levels.

ILF $= 0$ The sum of the contribution of the inelastic reactions will be made using a balance relationship involving the smooth cross sections.

An exception to this scheme is typically made when there is only one inelastic level within the unresolved energy range, because the flag can then just be set to its MT number and the special tabulation is not needed.

IOA The IOA is the other absorption flag for determining the contribution of “other absorptions” (no neutron out or destruction reactions).

IOA < 0 The “other absorption” cross section is zero within the entire unresolved resonance range.

IOA > 0 The value of IOA is a special MT number whose tabulation is the sum of the “other absorption” reactions.

IOA $= 0$ The sum of the contribution of the “other absorption” reactions will be made using a balanced relationship involving the smooth cross sections.

An exception to this scheme is typically made when there is only one “other absorption” reaction within the unresolved energy range, because the flag can then just be set to its MT number and the special tabulation is not needed.

Table 62: Possible values for the j argument.

j	Description
1	cumulative probability
2	total cross section/factor
3	elastic cross section/factor
4	fission cross section/factor
5	(n, γ) cross section/factor
6	neutron heating number/factor

IFF The IFF is the factors flag.

IFF = 0 The tabulations in the probability tables are cross sections.

IFF = 1 The tabulations in the probability tables are factors that must be multiplied by the corresponding “smooth” cross sections to obtain the actual cross sections.

The format of the **UNR Block** is given in Table 63. The $P(i, j, k)$ values, where

- $i = 1, \dots, N$,
- $j = 1, \dots, 6$,
- $k = 1, \dots, M$,

are what make up the probability tables. The argument j has special meaning depending on its value as shown in Table 62.

Table 63: UNR Block.

Location in XSS	Parameter	Description
JXS(23)	N	Number of incident energies where there is a probability table.
JXS(23)+1	M	Length of probability table.
JXS(23)+2	INT	Interpolation parameter between tables. [†]
JXS(23)+3	ILF	Inelastic competition flag.
JXS(23)+4	IOA	Other absorption flag.
JXS(23)+5	IFF	Factors flag.
JXS(23)+6	$E(i), i = 1, \dots, N$	Incident energies.
JXS(23)+6+N	$P(i, j, k)$	Probability tables.

[†] 2 linear-linear interpolation,
5 log-log interpolation

The ordering of the probability table entries, $P(i, j, k)$ is given in Table 64, which begins at PTABLE = JXS(23) + 6 + N .

Table 64: Order of probability table elements $P(i, j, k)$.

Location in XSS	Parameter	Description
Data for E(1)		
PTABLE	$\text{CDF}_1(l), l = 1, \dots, M$	Cumulative probabilities [†] for energy $i = 1$
PTABLE+ M	$\sigma_{t,1}(l), l = 1, \dots, M$	Total cross section/factors for energy $i = 1$
PTABLE+ $2M$	$\sigma_{s,1}(l), l = 1, \dots, M$	Elastic cross section/factors for energy $i = 1$
PTABLE+ $3M$	$\sigma_{f,1}(l), l = 1, \dots, M$	Fission cross section/factors for energy $i = 1$
PTABLE+ $4M$	$\sigma_{(n,\gamma),1}(l), l = 1, \dots, M$	(n, γ) cross section/factors for energy $i = 1$
PTABLE+ $5M$	$H_1(l), l = 1, \dots, M$	Heating number/factors for energy $i = 1$
Data for incident energy 2—same format for $E(1)$		
...		
Data for incident energy N—same format for $E(1)$		

[†] The cumulative probabilities are monotonically increasing from an implied (but not included) lower value of zero to the upper value of $P(i, 1, k = M) = 1.0$.

4.3.18 PTYPE Block

The **PTYPE Block** is the first of the particle production blocks used for neutron and charged particle production data. These particle production blocks are given only if the number of particles **NTYPE** = **NXS(7)** is different from zero. If the **PTYPE Block** is present, the **PTYPE Block** starts at the index **LTYPE** = **JXS(30)**.

Question: The particle type can be neutron, when is this actually used?

The **PTYPE Block** gives a list of particle types for which particle production data is available. This includes cross section data (given in the **SIGH Block**), angular distribution data (given in the **ANDH Block**) and secondary particle energy distribution data (given in the **DLWH Block**).

The format of the **PTYPE Block** is given in Table 65.

Table 65: PTYPE Block.

Location in XSS	Parameter	Description
LTYPE	IP_1	First particle type
LTYPE+1	IP_2	Second particle type
...		
LTYPE+NTYPE-1	IP_{NMT}	Last particle type

$\text{IP}_1, \dots, \text{IP}_{\text{NMT}}$ are particle identifiers given as follows:

- $\text{IP} = 1$ for neutrons
- $\text{IP} = 9$ for protons
- $\text{IP} = 31$ for deuterons

- IP = 32 for tritons
- IP = 33 for helions
- IP = 34 for aphas

4.3.19 NTRO Block

The [NTRO Block](#) gives the number of reactions for each of the particle types defined in the [PTYPE Block](#). If the [NTRO Block](#) is present, the [NTRO Block](#) starts at the index $LTYPE = JXS(31)$.

The format of the [NTRO Block](#) is given in Table 66.

Table 66: NTRO Block.

Location in XSS	Parameter	Description
LTYPE	NP_1	Number of reactions producing the first particle type
LTYPE+1	NP_2	Number of reactions producing the second particle type
...		
LTYPE+NTYPE-1	NP_{NMT}	Number of reactions producing the last particle type

4.3.20 IXS Block

The [IXS Block](#) gives 10 particle production locators for each of the particle types defined in the [PTYPE Block](#). If the [IXS Block](#) is present, the [IXS Block](#) starts at the index $NEXT = JXS(32)$. The [IXS Block](#) serves a similar function as the [JXS Array](#), in that it provides the locators to specific blocks of data, as laid out in Table 67.

Table 67: IXS array for particle type j , with $LTYPE = NEXT + 10NTYPE(j - 1)$ Block.

Location in XSS	Parameter	Description
LTYPE	HPD	Location of the total particle production and heating data
LTYPE+1	MTRH	Location of the particle production MT array
LTYPE+2	TYRH	Location of the particle production TYR data
LTYPE+3	LSIGH	Location of the particle production cross section locators
LTYPE+4	SIGH	Location of the particle production cross sections
LTYPE+5	LANDH	Location of the particle production angular distribution locators

Continued on next page

Table 67: IXS array for particle type j , with $LTYPE = NEXT + 10NTYPE(j - 1)$ Block (continued)

Location in XSS	Parameter	Description
LTYPE+6	ANDH	Location of the particle production angular distributions
LTYPE+7	LDLWH	Location of the particle production energy distribution locators
LTYPE+8	DLWH	Location of the particle production energy distributions
LTYPE+9	YH	Location of the particle production yield multipliers

With the exception of the [HPD Block](#), all other locators point to blocks similar to the ones already defined for neutrons and photons.

4.3.21 HPD Block

The [HPD Block](#) gives the total particle production cross section and the associated heating number for a given particle. If the particle production data is given (i.e. the $NTYPE \neq 0$, and the [PTYPE Block](#), the [NTRO Block](#) and [IXS Block](#) are given), the [HPD Block](#) is present. The [HPD Block](#) starts at the location index $HPD = XSS(())NEXT + 10 * (j - 1)$.

Table 68: HPD Block.

Location in XSS	Parameter	Description
HPD	IE	Energy grid index
HPD+1	N_E	Number of consecutive energies
HPD+2	$\sigma[E(K)],$	Total particle production cross section
HPD+2+NE	$E(l), l = 1, \dots, N_E$	Average heating numbers

5 Neutron Dosimetry

5.1 NXS Array

Table 69: NXS array element definitions for NXS ACE Table.

Element	Name	Description
1	—	Length of second block of data (XSS array)
2	ZA	$1000 * Z + A$
3	—	
4	NTR	Number of reactions
	...	
16	—	

5.2 JXS Array

Table 70: JXS array element definitions for JXS ACE Table.

Element	Name	Location	Description
1	LONE		First word of table
2	—		
3	MTR	MT	array
	...		
6	LSIG		Table of cross section locators
7	SIGD		Cross sections
	...		
22	END		Last word of this table
	...		
32	—		

6 Thermal Scattering $S(\alpha, \beta)$

Data from thermal $S(\alpha, \beta)$ tables provide a complete representation of thermal neutron scattering by molecules and crystalline solids. Cross sections for (coherent and incoherent) elastic and (incoherent) inelastic scattering are found on the tables. A coupled energy/angle representation is used to describe the spectra of inelastically scattered neutrons. Angular distributions for elastic scattering are also provided.

Four unique types of data blocks are associated with $S(\alpha, \beta)$ tables. We now briefly describe each of these four data block types and reference the sections in which their formats are detailed.

1. **ITIE Block**—contains the energy-dependent incoherent inelastic scattering cross sections. The **ITIE Block** always exists. See Section 6.3.1.
2. **ITCE Block** and **ITCEI Block**—contains the energy-dependent elastic scattering cross sections. The **ITCE Block** exists if the material has coherent and/or incoherent elastic scattering ($\text{NXS}(5) \neq 0$ and $\text{JXS}(4) \neq 0$). The **ITCEI Block** only exists for mixed mode elastic scattering ($\text{NXS}(5) = 5$ and $\text{JXS}(7) \neq 0$). See Section 6.3.3.
3. **ITXE Block**—contains coupled energy/angle distributions for incoherent inelastic scattering. The **ITXE Block** always exists. See Section 6.3.2.
4. **ITCA Block** and **ITCAI Block**—contains angular distributions for elastic scattering. The **ITCA Block** exists if the material has coherent and/or incoherent elastic scattering ($\text{NXS}(5) \neq 0$, $\text{JXS}(6) \neq 0$ and $\text{NXS}(6) \neq -1$). The **ITCAI Block** only exists for mixed mode elastic scattering ($\text{NXS}(5) = 5$, $\text{JXS}(9) \neq 0$ and $\text{NXS}(6) \neq -1$). See Section 6.3.4.

6.1 NXS Array

Table 71: NXS array element definitions for NXS ACE Table.

Element	Name	Description
1	—	Length of second block of data (XSS array)
2	IDPNI	Inelastic scattering mode
3	NIL	Inelastic dimensioning parameter
4	NIEB	Number of inelastic exiting energies
5	IDPNC	Elastic scattering mode (no elastic data=0, incoherent=3, coherent=4, mixed=5)
6	NCL	Elastic dimensioning parameter for the first elastic block
7	IFENG	Secondary energy mode (discrete=0, skewed=1, continuous=2)
8	NCLI	Elastic dimensioning parameter for the second elastic block
	...	
16	—	

6.2 JXS Array

Table 72: JXS array element definitions for JXS ACE Table.

Element	Name	Location	Description
1	ITIE		Inelastic energy table
2	ITIX		Inelastic cross sections
3	ITXE		Inelastic energy/angle distributions
4	ITCE		Elastic energy table (used for coherent elastic scattering if $\text{NXS}(5)=4$ or 5, and used for incoherent elastic scattering if $\text{NXS}(5)=3$)
5	ITCX		Elastic cross sections (used for coherent elastic scattering if $\text{NXS}(5)=4$ or 5, and used for incoherent elastic scattering if $\text{NXS}(5)=3$)
6	ITCA		Elastic angular distributions (used for coherent elastic scattering if $\text{NXS}(5)=4$ or 5, and used for incoherent elastic scattering if $\text{NXS}(5)=3$)
7	ITCEI		Elastic energy table (used for incoherent elastic scattering if $\text{NXS}(5)=5$)
8	ITCXI		Elastic cross sections (used for incoherent elastic scattering if $\text{NXS}(5)=5$)
9	ITCAI		Elastic angular distributions (used for incoherent elastic scattering if $\text{NXS}(5)=5$)
...			
32	—		

When a single mode of elastic scattering is used (either coherent when $\text{NXS}(5)=4$ or incoherent when $\text{NXS}(5)=3$), then only the first elastic block will be used. This is the way data was stored in the ACE format prior to the introduction of mixed mode elastic scattering in thermal scattering evaluations that combines both coherent and incoherent elastic scattering.

When using mixed mode elastic scattering (both coherent and incoherent elastic scattering are given, $\text{NXS}(5)=5$), the JXS array will contain an additional set of indices for a second elastic data block. In mixed mode, the first elastic blocks (pointed to by $\text{JXS}(4)$ to $\text{JXS}(6)$) are used for the coherent part and the second elastic blocks (pointed to by $\text{JXS}(7)$ to $\text{JXS}(9)$) are used for the incoherent part.

6.3 Format of Individual Data Blocks

6.3.1 ITIE Block

The format of the [ITIE Block](#) is given in Table 73. The index at the start of the block is $S_{\text{ITIE}}=\text{JXS}(1)$. Note that $\text{JXS}(2)=\text{JXS}(1)+1+N_{in}$. Linear-linear interpolation is assumed between adjacent energies.

Table 73: ITIE Block.

Location in XSS	Parameter	Description
S_{ITIE}	N_{in}	Number of inelastic energies

Continued on next page

Table 73: ITIE Block (continued)

Location in XSS	Parameter	Description
$S_{\text{ITIE}}+1$	$E_{in}(l), l = 1, \dots, N_{in}$	Energies
$S_{\text{ITIE}}+1+N_{in}$	$\sigma_{in}(l), l = 1, \dots, N_{in}$	Inelastic cross sections

6.3.2 ITXE Block

The format of the coupled energy/angle distribution for incoherent inelastic scattering is governed by the value of $\text{NXS}(7)$. There are three possibilities:

$\text{NXS}(7) = 0$ equally-likely discrete cosines and energies (Table 74)

$\text{NXS}(7) = 1$ skewed distribution of discrete cosines and energies (Table 74)

$\text{NXS}(7) = 2$ continuous distribution of outgoing energies and equally-likely discrete cosines (Table 75 and Table 76)

The format of the **ITXE Block** for $\text{NXS}(7) < 2$ is given in Table 74. The index at the start of the block is $S_{\text{ITXE}} = \text{JXS}(3)$. For each incident energy from the **ITIE Block**, $N' = \text{NXS}(4)$ discrete outgoing energies are given. For each pair of incident and outgoing energies, $N_\mu = \text{NXS}(3) + 1$ discrete cosines are given. The incident inelastic energy grid $E_{in}(l)$ is given in the **ITIE Block**, and linear-linear interpolation is assumed between adjacent values of E_{in} .

Table 74: **ITXE Block** for $\text{NXS}(7) < 2$.

Location in XSS	Parameter	Description
S_{ITXE}	$E_1^{\text{out}}[E_{in}(1)]$	First of NIEB [†] outgoing energies for inelastic scattering at $E_{in}(1)$
$S_{\text{ITXE}}+1$	$\mu_l(1 \rightarrow 1), l = 1, \dots, N_\mu$	Discrete cosines for scattering from $E_{in}(1)$ to $E_1^{\text{out}}[E_{in}(1)]$
$S_{\text{ITXE}}+1+N_\mu$	$E_2^{\text{out}}[E_{in}(1)]$	Second of NIEB outgoing energies for inelastic scattering at $E_{in}(1)$
$S_{\text{ITXE}}+2+N_\mu$	$\mu_l(1 \rightarrow 2), l = 1, \dots, N_\mu$	Discrete cosines for scattering from $E_{in}(1)$ to $E_2^{\text{out}}[E_{in}(1)]$
$\vdots$	$\vdots$	$\vdots$
$S_{\text{ITXE}}+(N'-1)(1+N_\mu)$	$E_{N'}^{\text{out}}[E_{in}(1)]$	Last of NIEB outgoing energies for inelastic scattering at $E_{in}(1)$
$S_{\text{ITXE}}+(N'-1)(1+N_\mu)+1$	$\mu_l(1 \rightarrow N'), l = 1, \dots, N_\mu$	Discrete cosines for scattering from $E_{in}(1)$ to $E_{N'}^{\text{out}}[E_{in}(1)]$
(Repeat for all remaining values of E_{in})		

[†] The number of outgoing energies **NIEB** is determined as $\text{NXS}(4)$.

When $\text{NXS}(7) = 0$, each of the $\text{NXS}(4)$ discrete outgoing energies for a given incident energy are

equally probable. When $\text{NXS}(7) = 1$, the selection of the discrete outgoing energies is skewed such that the first two and last two outgoing energies have a lower probability of being selected than all other outgoing energies. The first and last energies have a relative probability of 1, the second and second-to-last energies have a relative probability of 4, and all other energies have a relative probability of 10.

Because the use of discrete outgoing energies and cosines can result in unphysical spikes in the neutron flux spectrum at thermal energies, some Monte Carlo codes attempt to “smear” the outgoing energies and cosines to produce a smoother distribution (that more closely approximates a continuous distribution with $\text{NXS}(7) = 2$).

When $\text{NXS}(7) = 2$, the distribution of outgoing energies for each incident energy is continuous in energy and specified by a probability density function and cumulative distribution function. The format of the [ITXE Block](#) in this case is given in Table 75 and Table 76. As before, the index at the start of the block is $S_{\text{ITXE}} = \text{JXS}(3)$. Unlike in the $\text{NXS}(7) < 2$ cases, the number of outgoing energies for each incident energy is allowed to vary. The number of discrete cosines, $N_\mu = \text{NXS}(3) - 1$, remains the same for each pair of incident and outgoing energies, however.

Table 75: [ITXE Block](#) for $\text{NXS}(7) = 2$.

Location in XSS	Parameter	Description
S_{ITXE}	$L(l), l = 1, \dots, N_{in}^\dagger$	Location in XSS of distribution for incident energy l
$S_{\text{ITXE}} + N_{in}$	$N'(l), l = 1, \dots, N_{in}$	Number of outgoing energies for incident energy l

[†] The number of incoming energies N_{in} for incoherent inelastic scattering is given in the [ITIE Block](#).

Table 76: [ITXE Block](#) for $\text{NXS}(7) = 2$ (continued).

Location in XSS	Parameter	Description
$L(1)+1$	$E_1^{\text{out}}[E_{in}(1)]$	First of NIEB outgoing energies for inelastic scattering at $E_{in}(1)$
$L(1)+2$	$\text{PDF}_1[E_{in}(1)]$	Probability density function value for $E_1^{\text{out}}[E_{in}(1)]$
$L(1)+3$	$\text{CDF}_1[E_{in}(1)]$	Cumulative distribution function value for $E_1^{\text{out}}[E_{in}(1)]$
$L(1)+4$	$\mu_l(1 \rightarrow 1), l = 1, \dots, N_\mu$	Discrete cosines for scattering from $E_{in}(1)$ to $E_1^{\text{out}}[E_{in}(1)]$
$L(1)+4+N_\mu$	$E_2^{\text{out}}[E_{in}(1)]$	Second of NIEB outgoing energies for inelastic scattering at $E_{in}(1)$

Continued on next page

Table 76: [ITXE Block](#) for $\text{NXS}(7) = 2$ (continued) (continued)

Location in XSS	Parameter	Description
$\text{L}(1)+5+N_\mu$	$\text{PDF}_2[E_{in}(1)]$	Probability density function value for $E_2^{out}[E_{in}(1)]$
$\text{L}(1)+6+N_\mu$	$\text{CDF}_2[E_{in}(1)]$	Cumulative distribution function value for $E_2^{out}[E_{in}(1)]$
$\text{L}(1)+7+N_\mu$	$\mu_l(1 \rightarrow 2), l = 1, \dots, N_\mu$	Discrete cosines for scattering from $E_{in}(1)$ to $E_2^{out}[E_{in}(1)]$
$\vdots$	$\vdots$	$\vdots$
$\text{L}(1)+1+(N'(1)-1)(3+N_\mu)$	$E_{N'(1)}^{out}[E_{in}(1)]$	Last of $N'(1)$ outgoing energies for inelastic scattering at $E_{in}(1)$
$\text{L}(1)+2+(N'(1)-1)(3+N_\mu)$	$\text{PDF}_{N'(1)}[E_{in}(1)]$	Probability density function value for $E_{N'(1)}^{out}[E_{in}(1)]$
$\text{L}(1)+3+(N'(1)-1)(3+N_\mu)$	$\text{CDF}_{N'(1)}[E_{in}(1)]$	Cumulative distribution function value for $E_{N'(1)}^{out}[E_{in}(1)]$
$\text{L}(1)+4+(N'(1)-1)(3+N_\mu)$	$\mu_l(1 \rightarrow N'(1)), l = 1, \dots, N_\mu$	Discrete cosines for scattering from $E_{in}(1)$ to $E_{N'(1)}^{out}[E_{in}(1)]$
(Repeat for all remaining values of E_{in})		

6.3.3 ITCE Block

The format of the [ITCE Block](#) and [ITCEI Block](#) is given in Table 77. The index at the start of the [ITCE Block](#) and [ITCEI Block](#) is respectively $S_{\text{ITCE}}=\text{JXS}(4)$ and $S_{\text{ITCEI}}=\text{JXS}(7)$.

Table 77: ITCE Block.

Location in XSS	Parameter	Description
S_{ITCE}	N_{el}	Number of elastic energies
$S_{\text{ITCE}}+1$	$E_{el}(l), l = 1, \dots, N_{el}$	Energies
$S_{\text{ITCE}}+1+N_{el}$	$P(l), l = 1, \dots, N_{el}$	(See below)

For incoherent elastic scattering (stored in [ITCE Block](#) if $\text{NXS}(5) = 4$ and stored in [ITCEI Block](#) if $\text{NXS}(5) = 5$),

$$P(l) = \sigma_{el}(E_{el}(l)) \quad (22)$$

with linear-linear interpolation between points. For coherent elastic scattering (stored in [ITCE Block](#) if $\text{NXS}(5) = 4$ or 5),

$$P(l) = E \cdot \sigma_{el}(E), \quad E_{el}(l) \leq E < E_{el}(l+1). \quad (23)$$

In this case, the energies $E_{el}(l)$ correspond to Bragg edges, and between two energies the cross section is determined by inverting Equation 23:

$$\sigma_{el}(l) = \frac{P(l)}{E}, \quad E_{el}(l) \leq E < E_{el}(l+1). \quad (24)$$

Also note that $\sigma_{el}(E) = 0$ below $E_{el}(1)$. However, above $E_{el}(N_{el})$, $\sigma_{el}(E) = P(N_{el})/E$.

6.3.4 ITCA Block

The format of the [ITCA Block](#) and [ITCAI Block](#) is given in Table 78. The index at the start of the [ITCA Block](#) and [ITCAI Block](#) is respectively $S_{\text{ITCA}}=\text{JXS}(6)$ and $S_{\text{ITCA}}=\text{JXS}(9)$. For each incident energy from the [ITCE Block](#) and [ITCEI Block](#) respectively, $N_\mu = \text{NXS}(6)+1$ and $N_\mu = \text{NXS}(8)+1$ discrete cosines are given.

Table 78: ITCA Block.

Location in XSS	Parameter	Description
S_{ITCA}	$\mu_l[E_{el}(1)], l = 1, \dots, N_\mu$	Discrete cosines for elastic scattering at $E_{el}(1)$
$S_{\text{ITCA}}+N_\mu$	$\mu_l[E_{el}(2)], l = 1, \dots, N_\mu$	Discrete cosines for elastic scattering at $E_{el}(2)$
$\vdots$	$\vdots$	$\vdots$
$S_{\text{ITCA}}+(N_{el}-1)N_\mu$	$\mu_l[E_{el}(N_{el})], l = 1, \dots, N_\mu$	Discrete cosines for elastic scattering at $E_{el}(N_{el})$

The incident elastic energy grid $E_{el}(l)$ is given in the [ITCE Block](#) or [ITCEI Block](#). Linear-linear interpolation is assumed between adjacent values of E_{el} .

7 Continuous-Energy Photon

7.1 NXS Array

Table 79: NXS array element definitions for NXS ACE Table.

Element	Name	Description
1	—	Length of second block of data (XSS array)
2	Z	Atomic number
3	NES	Number of energies
4	NFLC	Length of the fluorescence data divided by 4
5	NSH	Number of electron shells
	...	
16	—	

7.2 JXS Array

Table 80: JXS array element definitions for JXS ACE Table.

Element	Name	Location Description
1	ESZG	Energy table
2	JINC	Incoherent form factors
3	JCOH	Coherent form factors
4	JFLO	Fluorescence data
5	LHNM	Heating numbers
6	LNEPS	Number of electrons per shell
7	LBEPS	Binding energy per shell
8	LPIPS	Probability of interaction per shell
9	LSWD	Array of offsets to the shell-wise data
10	SWD	Shell-wise data in PDF and CDF form
	...	
32	—	